

<https://helda.helsinki.fi>

---

## The current state of CO<sub>2</sub> flux chamber studies in the Arctic tundra : A review

Virkkala, Anna-Maria

2018-04

---

Virkkala , A-M , Virtanen , T , Lehtonen , A , Rinne , J & Luoto , M 2018 , ' The current state of CO<sub>2</sub> flux chamber studies in the Arctic tundra : A review ' , Progress in Physical Geography , vol. 42 , no. 2 , pp. 162-184 . <https://doi.org/10.1177/0309133317745784>

---

<http://hdl.handle.net/10138/307190>

<https://doi.org/10.1177/0309133317745784>

---

cc\_by\_nc\_nd

acceptedVersion

---

*Downloaded from Helda, University of Helsinki institutional repository.*

*This is an electronic reprint of the original article.*

*This reprint may differ from the original in pagination and typographic detail.*

*Please cite the original version.*

**The current state of CO<sub>2</sub> flux chamber studies in the tundra:  
a review**

Journal:	<i>Progress in Physical Geography</i>
Manuscript ID	PPG-17-050.R1
Manuscript Type:	Main Article
Keywords:	Arctic, chamber, ecosystem respiration, gross primary production, net ecosystem exchange, tundra
Abstract:	<p>The Arctic tundra plays an important role in the carbon cycle as it stores 50 % of global soil organic carbon reservoirs. The processes (fluxes) regulating these stocks are predicted to change due to direct and indirect effects of climate change. Understanding the current and future carbon balance calls for a summary of the level of knowledge regarding chamber-derived CO<sub>2</sub> flux studies. Here, we describe progress from recently (2000-2016) published studies of growing-season CO<sub>2</sub> flux chamber measurements, namely GPP (gross primary production), ER (ecosystem respiration), and NEE (net ecosystem exchange), in the tundra region. We review the study areas and designs along with the explanatory environmental drivers used. Most of the studies were conducted in Alaska and Fennoscandia, and we stress the need for measuring fluxes in other tundra regions, particularly in more extreme climatic, productivity, and soil conditions. Soil respiration and other greenhouse gas measurements were seldom included in the studies. Although most of the environmental drivers of CO<sub>2</sub> fluxes have been relatively well investigated (such as the effect of vegetation type and soil microclimate on fluxes), soil nutrients, other greenhouse gases and disturbance regimes require more research as they define the future carbon balance. Particular attention should be paid to the effects of shrubification, geomorphology, and other disturbance effects such as fire events, and disease and herbivore outbreaks. An improved conceptual framework and understanding of underlying processes of biosphere-atmosphere CO<sub>2</sub> exchange will provide more information on carbon cycling in the tundra.</p>

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1

2 The current state of CO<sub>2</sub> flux chamber studies in the Arctic tundra: a review

3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

3

4

5 **Abstract**

6 The Arctic tundra plays an important role in the carbon cycle as it stores 50 % of global soil  
7 organic carbon reservoirs. The processes (fluxes) regulating these stocks are predicted to  
8 change due to direct and indirect effects of climate change. Understanding the current and  
9 future carbon balance calls for a summary of the level of knowledge regarding chamber-  
10 derived CO<sub>2</sub> flux studies. Here, we describe progress from recently (2000-2016) published  
11 studies of growing-season CO<sub>2</sub> flux chamber measurements, namely GPP (gross primary  
12 production), ER (ecosystem respiration), and NEE (net ecosystem exchange), in the tundra  
13 region. We review the study areas and designs along with the explanatory environmental  
14 drivers used. Most of the studies were conducted in Alaska and Fennoscandia, and we  
15 stress the need for measuring fluxes in other tundra regions, particularly in more extreme  
16 climatic, productivity, and soil conditions. Soil respiration and other greenhouse gas  
17 measurements were seldom included in the studies. Although most of the environmental  
18 drivers of CO<sub>2</sub> fluxes have been relatively well investigated (such as the effect of vegetation  
19 type and soil microclimate on fluxes), soil nutrients, other greenhouse gases and  
20 disturbance regimes require more research as they might define the future carbon balance.  
21 Particular attention should be paid to the effects of shrubification, geomorphology, and  
22 other disturbance effects such as fire events, and disease and herbivore outbreaks. An  
23 improved conceptual framework and understanding of underlying processes of biosphere-  
24 atmosphere CO<sub>2</sub> exchange will provide more information on carbon cycling in the tundra.

25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

25

26 **Keywords**

27 Arctic, tundra, chamber, net ecosystem exchange, gross primary production, ecosystem  
28 respiration

## I Introduction

Tundra regions are expected to experience stronger climate warming than other regions in the world (Hinzman et al., 2013; Stocker et al., 2013). They are also highly sensitive to environmental change, which may potentially alter their ecological dynamics (Hinzman et al., 2005; McGuire et al., 2009; Post et al., 2009). Higher air temperatures have already been observed to alter permafrost extent (Osterkamp and Romanovsky, 1999; Schuur et al., 2009), vegetation distribution (Elmendorf et al., 2012; Forbes et al., 2010; Hudson and Henry, 2009; Tape et al., 2006), and hydrological cycles (Mernild et al., 2012; Overeem and Syvitski, 2010; Smol and Douglas, 2007). These changes are tightly linked to the tundra carbon cycle which plays an important role in the global carbon balance: northern tundra regions store approximately 50 % of global soil organic carbon (SOC) stocks (Hugelius et al., 2014; McGuire et al., 2009; Tarnocai et al., 2009). Whether tundra regions will act as a carbon sink, or as a carbon source, in a warmer climate remains uncertain (Belshe et al., 2013; McGuire et al., 2012; Sitch et al., 2007).

The SOC stocks and the carbon balance are regulated mainly by CO<sub>2</sub> and CH<sub>4</sub> exchange between the biosphere and the atmosphere (McGuire et al., 2010a). Moreover, the lateral transport of organic and inorganic carbon in soil water transfers carbon to streams and rivers in a dissolved or particulate form (McGuire et al., 2010b; McGuire et al., 2009; Parmentier et al., 2017). Two processes, or fluxes, dominate the biosphere-atmosphere exchange: CO<sub>2</sub> uptake by ecosystems via photosynthesis (GPP: gross primary production), and carbon release to the atmosphere via plant and microbial respiratory losses (ER: ecosystem respiration) (Callaghan et al., 2004). The CO<sub>2</sub> balance between these two fluxes is called NEE (net ecosystem exchange).

Future climate warming can affect CO<sub>2</sub> fluxes in multiple ways (Arens et al., 2008; Biasi et al., 2008; Bokhorst et al., 2010). On the one hand, warming might accelerate soil respiration by heterotrophs, i.e. the microbial decomposition of organic matter, thus increasing respiratory carbon losses (Grogan and Chapin, 2000; Nadelhoffer et al., 1991). These emissions may be multiplied if permafrost soils thaw and the carbon stored therein will

1  
2  
3 70 become available for decomposition (Kuhry et al., 2010; van Huissteden and Dolman,  
4 71 2012). On the other hand, increases in temperature, greater release of available soil  
5 72 nutrients via accelerated decomposition, and a longer growing season have the potential  
6 73 to stimulate GPP and strengthen the carbon sink (Cahoon et al., 2012; Leffler et al., 2016).  
7 74 Changes in the CO<sub>2</sub> fluxes can therefore either increase or decrease NEE and thus create  
8 75 either a negative or positive feedback to global warming. A recent meta-analysis based on  
9 76 54 studies suggested that there has been an increase in the CO<sub>2</sub> source due to higher ER,  
10 77 resulting in amplified carbon losses to the atmosphere in warmer conditions (Belshe et al.,  
11 78 2013).

12  
13  
14  
15  
16 79  
17  
18 80 Though climatically controlled, the magnitudes, directions, and feedbacks of GPP and ER  
19 81 are ecosystem- and scale-dependent (Oberbauer et al., 2007; Pare and Bedard-Haughn,  
20 82 2012). Tundra landscape heterogeneity (Fletcher et al., 2012; Post et al., 2009; Virtanen and  
21 83 Ek, 2014) is apparent at multiple scales. At the landscape scale, diverse environmental  
22 84 conditions and vegetation types create mosaic of ecosystems, for example, bogs, and  
23 85 barrens. These ecosystems, in turn, are made up of diverse plant-communities, such as  
24 86 heaths, tussocks, or hummocks. At this fine scale, environmental conditions can change  
25 87 within only a few meters (Aalto et al., 2013). Such variation might offset climatological  
26 88 effects and buffer climate change -induced modifications in CO<sub>2</sub> fluxes (Post et al., 2009).  
27 89 For example, the environmental gradients of soil moisture are an important driver of CO<sub>2</sub>  
28 90 fluxes (Nobrega and Grogan, 2008; Oberbauer et al., 2007; Sharp et al., 2013). However, the  
29 91 fine-scale variation of CO<sub>2</sub> fluxes is not yet fully understood. This may hinder our ability to  
30 92 predict the future carbon balance.

31  
32  
33  
34  
35  
36  
37 93  
38  
39 94 GPP, ER, and NEE are studied with various methods in the field. There are two widely used  
40 95 techniques to measure CO<sub>2</sub> fluxes between ecosystems and the atmosphere: the  
41 96 micrometeorological eddy covariance (EC) technique, and chamber measurement  
42 97 techniques. EC measures fluxes continuously at an ecosystem scale in areas between 100  
43 98 m<sup>2</sup> to 10 000 m<sup>2</sup>, thus averaging over fine-scale heterogeneity (Kade et al., 2012;  
44 99 Maanavilja et al., 2011). However, using detailed land cover and wind direction information,  
45 100 flux sources can to some extent be estimated by footprint modelling techniques (Fox et al.,  
46 101 2008; Soegaard et al., 2000). Studies investigating individual plant communities and fine-  
47 102 scale spatial variability (100–10 000 cm<sup>2</sup>) in the tundra are usually conducted using  
48 103 chambers. Both abovementioned methods have their limitations. The costs of setting up  
49 104 and maintaining an EC tower are higher than for the chamber method. Additionally, EC  
50 105 requires more electricity, technical knowledge, and understanding of surface-layer  
51 106 meteorology; whereas chambers are widely used due to low power consumption and  
52 107 simple operation. However, manual chamber measurements are laborious. With the

chamber method, the different components of CO<sub>2</sub> exchange can be studied, thus providing important data on the processes regulating biosphere-atmosphere interactions (McGuire et al., 2012).

In order to understand CO<sub>2</sub> fluxes at multiple scales, reviews and meta-analyses are important to summarize the current knowledge. Comprehensive works exist on the Arctic carbon cycle as a whole (McGuire et al., 2009; Parmentier et al., 2017), on the vulnerability of permafrost carbon (Abbott et al., 2016; Ping et al., 2015; Schuur et al., 2008), and on the CO<sub>2</sub> and carbon balance of tundra regions during the past 20-40 years (Belshe et al., 2013; McGuire et al., 2010b, McGuire et al., 2012, Oechel et al., 1993, Oechel et al., 2000). Although these analyses include chamber studies, the main focus has been on the effects of large-scale drivers and temporal changes on CO<sub>2</sub> fluxes. To our knowledge, no previous reviews or meta-analyses focus on chamber studies of GPP, ER, and NEE with an emphasis on fine-scale vegetation and soil variables.

In this review, we synthesize the progress of chamber-derived growing season CO<sub>2</sub> flux studies in terrestrial Arctic tundra. We focus on chamber studies as they have been and will continue to be a central and cost-efficient method to study the underlying processes in CO<sub>2</sub> exchange, and are able to account for the fine-scale spatial variability of these processes. GPP and ER are the most important fluxes considered (Cahoon et al., 2012; Illeris et al., 2004), as they are processes that govern the net exchange of CO<sub>2</sub> between an ecosystem and the atmosphere. Furthermore, the balance of these two, NEE, is also incorporated in the review. We review the study areas and study designs where GPP, ER, and NEE are investigated and identify environmental conditions not sufficiently covered by these studies. Our aim is to examine the explanatory variables used and whether they correspond to mechanisms of known importance. Chapter 2 will give a short introduction to the tundra carbon cycle and chapters 3 to 6 encompass a systematic review of 93 articles found via ISI Web of Science (WoS) and subsequently from the references therein. We include recent publications only (studies published 2000 – 2016: Supplementary Material 1, S1) to determine the current state of CO<sub>2</sub> flux chamber studies in the tundra environment.

## II Carbon cycle in a tundra environment

The spatial extent of Arctic tundra regions (from here on, tundra regions) can be defined in various ways. Here, we consider the tundra as a biome of treeless vegetation and their adjacent tree-line areas in the arctic and oro-arctic regions (as suggested by Virtanen et al.

2016). Sub-Arctic and Arctic peatlands were included in the review whilst boreal regions were excluded as we concentrated on tundra patterns and processes only. Thus defined, tundra regions cover 8.7 % of global terrestrial areas (Dinerstein et al., 2017) and are characterized by low summer temperatures and long winters (Billings and Mooney, 1968).

Due to low solar energy input and harsh climatic seasonality, primary production, respiration, and decomposition rates are slow. Significant seasonal cycles in these processes exist largely due to snow cover and the short, two-to-four-month growing season (Leffler et al., 2016; van der Molen et al., 2007). Photosynthesis only occurs during the summer, whereas decomposition and respiration occur throughout the year, albeit at slower rates in winter (Larsen et al., 2007). The growing season is the most active season for GPP and ER, although spring and autumn CO<sub>2</sub> fluxes can contribute up to 19 % of annual GPP (Larsen et al. 2007). GPP and ER may fluctuate on inter-annual and longer time scales due to the variations in for example, climatic parameters. Diurnal cycles are also notable: daytime GPP usually offsets ER, but at night ER becomes dominant (López-Blanco et al., 2016). NEE, defined as NEE=ER-GPP, is usually a small difference between two large fluxes which makes it sensitive to small changes. Based on observations, tundra regions can be either carbon sources or sinks with large temporal variation on inter-annual and decadal scales (Lopez-Blanco et al., 2016). Typical NEE values are, for example, -79 - -41 g C m<sup>-2</sup> yr<sup>-1</sup> in a Russian erect shrub tundra - wetland complex (Marushchak et al., 2013), -4 g C m<sup>-2</sup> yr<sup>-1</sup> in a Fennoscandian fen (Aurela et al., 2004), and -2.3 g C m<sup>-2</sup> yr<sup>-1</sup> in an Arctic valley in Greenland (Soegaard et al., 2000), with negative values indicating a net carbon uptake by ecosystems. These estimates are smaller than for the neighboring boreal and temperate regions, but they have the potential to change in a warmer climate (Parker et al. 2015; Hartley et al. 2012).

Although tundra regions lack the trees that dominate carbon exchange in forested ecosystems, they have highly variable vegetation patterns. One of the most common ways to describe tundra vegetation is the Circumpolar Arctic Vegetation Map (CAVM; e.g. Pearson et al., 2013; Reynolds et al., 2008) created by Walker et al., (2005). They divide the Arctic based on the general appearance of vegetation into the following habitat-scale physiognomic categories and further into specific mapping units based on plant functional types (PFT's): barren (B), graminoid (G), prostrate-shrub (P), erect-shrub (S) and wetland (W) tundra, all of which have distinct CO<sub>2</sub> flux patterns (Figure 1). GPP usually exceeds ER, leading to a net uptake of carbon (negative NEE) during the growing season in all CAVM categories except in barren vegetation types. In general, deciduous shrubs and graminoids have the highest GPP due to high leaf area and vegetation cover, with graminoids having even larger ER due to their faster metabolism (Cahoon et al., 2012; Hobbie, 1996; Nobrega



and Grogan, 2008; Oberbauer et al., 2007). Yet, the annual leaf development cycle of deciduous plants shortens their carbon uptake period in the early and late growing season. Evergreen shrubs and cryptograms, on the other hand, start photosynthesizing as soon as the ground is snow-free (Douma et al., 2007; Street et al., 2012).

### III Study designs and locations

#### 1 Overview of the studies

During the 21<sup>st</sup> century, most of the studies have focused on the effects of climate warming and related feedbacks on CO<sub>2</sub> fluxes (Biasi et al., 2008; Boelman et al., 2003; Welker et al., 2004), differences between vegetation types (Shaver et al., 2007; Williams et al., 2006), hydrological effects (Dagg and Lafleur, 2011; Nobrega and Grogan, 2008), and growing season dynamics (Bäckstrand et al., 2008; Larsen et al., 2007). The effect of permafrost thaw on CO<sub>2</sub> fluxes has not received a lot of attention until lately (Natali et al., 2011; Schuur et al., 2009b; Vogel et al., 2009). The most cited articles (>100 citations) within the literature reviewed focused on the effects of experimental warming and permafrost thaw on CO<sub>2</sub> fluxes and their spatial variation (S3).

The geographical distribution of observations in the CO<sub>2</sub> flux chamber studies is clustered in Alaska (32 out of 93 studies) and Fennoscandia (27), with smaller clusters scattered across Greenland (18), Russia (13), and northern Canada and Svalbard (8, Figure 2A). The observation sites do not cover the environmental variation, nor the extreme conditions, found in the Arctic (Figure 2B). Extreme conditions are generally found in the middle and eastern parts of Russia, and along coastlines and islands of northern Canada and Greenland. The environmental conditions in Alaska and Fennoscandia are characterized by relatively high precipitation, semi-oceanic climates, and acidic or mildly acidic soil conditions. In addition to these spatial limitations, continuous sampling of CO<sub>2</sub> fluxes remains a challenge (McGuire et al., 2012). While there are many studies with measurements from multiple growing seasons (Arens et al., 2008; Olivas et al., 2010), there are not many year-round observations (S4). The cold season (autumn, winter and spring) has been included in some studies, mainly in order to understand the role of non-growing season vs. growing season CO<sub>2</sub> fluxes and their drivers (Grogan and Jonasson, 2005; Kade et al., 2012; Larsen et al., 2007).



1  
2  
3 218 CO<sub>2</sub> flux chamber studies are usually based on relatively small data sets (median number of  
4 219 sites studied four). With small data sets, all environmental gradients (e.g. topography,  
5 220 radiation, hydrology, productivity) are difficult to cover. Often, studies are designed to  
6 221 simplify and understand the variation with the help of different classifications mainly based  
7 222 on vegetation (Christensen et al., 2000 on Cassiope, hummock, continuous fen, grassland  
8 223 and *Salix arctica*), soil moisture (Welker et al., 2000 on dry and wet), or a combination of  
9 224 these two (Dagg and Lafleur, 2010 on dry birch, mesic birch, dry tussock, wet tussock, dry  
10 225 heath, mesic heath, wet sedge). The topographic gradient has been considered mostly in  
11 226 the peatland studies with microtopographically varying hummocks and hollows (e.g.  
12 227 Poyatos et al., 2014), though there are some coarser scale topo-sequence studies (Williams  
13 228 et al., 2006).

14  
15  
16  
17  
18  
19 229  
20  
21 230 Based on our findings, future research should focus more on extreme conditions. Areas of  
22 231 high precipitation and low temperatures (e.g. southeastern parts of Greenland) or high  
23 232 temperatures (e.g. southern Alaska) should also receive further attention. Additionally,  
24 233 neutral conditions (topsoil pH>6, e.g. along the coastlines of Nunavut and Siberia) and  
25 234 higher elevations (e.g. in eastern Siberia) need to be considered. Relatively large SOC  
26 235 stocks are found in the Northwestern or Yukon Territories in Canada, which have not  
27 236 received much consideration. As fluxes in Alaska and Fennoscandia are the most studied  
28 237 and cited, it seems that our understanding of CO<sub>2</sub> fluxes is limited to these regions and  
29 238 important knowledge regarding for example High-Arctic environments has not been  
30 239 adequately considered. We realize that logistical and infrastructural reasons are, to some  
31 240 extent, responsible for this pattern, but this may potentially lead to misinterpretations of  
32 241 the CO<sub>2</sub> exchange of the whole tundra region. Moreover, studies in calcareous regions are  
33 242 needed as the higher availability of nutrients in neutral areas may cause CO<sub>2</sub> fluxes to vary  
34 243 more than in acidic landscapes with low nutrient availability (Williams et al., 2006).  
35 244 Furthermore, we emphasize the need for better observational studies to understand the  
36 245 spatio-temporal variation of CO<sub>2</sub> fluxes along environmental gradients as the drivers and  
37 246 dynamics might change due to global warming (Shaver, et al., 2013; Zamołodchikov, 2015).  
38 247 We encourage future studies to measure CO<sub>2</sub> fluxes from various topographical conditions  
39 248 from fine, 1–10 m (Scherrer and Körner, 2011), to ecosystem and landscape scales, 10–1000  
40 249 m (Billings, 1973), as the response of biota to climate warming varies depending on  
41 250 topographic position.

42  
43  
44 251  
45  
46 252 *2 CO<sub>2</sub> flux chamber measurement methods*

47  
48  
49 253

Various chamber systems have been used for studying CO<sub>2</sub> fluxes. Most of the tundra CO<sub>2</sub> flux studies were conducted with manual chambers (82 studies) while automated systems are still rare (11) because of the high costs and infrastructure challenges (S4; e.g. Griffis et al., 2000; López-Blanco et al., 2016; Natali et al., 2011; Poyatos et al., 2014).

Chamber size plays an important role when studying CO<sub>2</sub> fluxes. Tundra regions are microtopographically heterogeneous and, therefore, smaller chambers (<30 cm in diameter) are more practical. However, with a larger chamber, the erroneous influence of chamber edges on measurements decreases. Based on our literature survey, chamber sizes (when measured from edges or diameter of the chamber) and heights vary from 15 to 100 cm. The average height is relatively low, limiting the vegetation type studied (S4). Regardless of chamber type and size, the air-proof sealing of the chamber into the ground is an important step when measuring fluxes (Hutchinson and Livingston, 2001). Traditionally, plastic or aluminium collars are put into the ground (depth 2-10 cm) and chamber placed atop of them. Collars are positioned into the ground just minutes, days (Dagg and Lafleur, 2011; Stewart et al., 2013) or weeks before the measurement (Nykanen et al. 2003; Heikkinen et al. 2002), or in the preceding year (Grogan and Jonasson, 2005). However, placing the collars into rocky soils or soils with long horizontal aboveground shoots and roots typical to some tundra plants (e.g. *Empetrum hermaphroditum*, *Dryas* sp.) without cutting them is a challenge. A method that uses the combination of a plastic skirt and metal chains instead of, or in addition to, collars has been developed and is widely used in the studies (Street et al., 2012; Street et al., 2007).

We stress the need for long-term measurements of CO<sub>2</sub> fluxes with automated chambers, as they provide important information about local carbon cycle dynamics and episodic events in the Arctic. Moreover, we recommend researchers to carefully consider the chamber size used, and the timing and placement of the collar into the ground. Measurement devices and calculation techniques also have an impact on the measured fluxes. For example, CO<sub>2</sub> fluxes may be underestimated when using linear regression as compared to nonlinear methods (Kutzbach et al., 2007).

### 3 Studied fluxes

From the global carbon balance perspective, NEE is an important flux variable to study as it describes whether the environment is acting as a CO<sub>2</sub> sink or source (Belshe et al., 2013). Yet, NEE itself is not a process but a balance between two simultaneous processes, GPP and ER, that are more strongly linked to environmental drivers. Although studying NEE is

crucial, many articles state that the future carbon balance is regulated by ER (Cahoon et al., 2012; Biasi et al., 2008) or soil respiration (Knowles et al., 2016; Luo et al., 2016). Despite this, soil respiration was measured in relatively few studies (15) together with GPP, ER, and NEE. Moreover, CO<sub>2</sub> fluxes are linked to other biogeochemical (N<sub>2</sub>O, CH<sub>4</sub> and H<sub>2</sub>O) fluxes. Methane (CH<sub>4</sub>) fluxes were studied in 24 studies, out of which most of the measurements were from wetlands (S4). Other fluxes, such as nitrous oxide (N<sub>2</sub>O; e.g. Brummell et al., 2012; Kelsey et al., 2016) or water (H<sub>2</sub>O; e.g. Douma et al., 2007) were studied even less in CO<sub>2</sub> flux chamber studies, partly because according to current knowledge, N<sub>2</sub>O emissions are significant in only non-vegetated patches with high soil organic matter content (Marushchak et al., 2011). They are increasingly measured alone (Flessa et al., 2008; Gil et al., 2017; Voigt et al., 2017), without CO<sub>2</sub> exchange measurements. Chamber-derived CO<sub>2</sub> fluxes have also been investigated together with EC and leaf cuvette methods (Heikkinen et al. 2002; Vourlitis et al. 2000). Studies with EC aimed to 1) describe the annual variation of CO<sub>2</sub> fluxes more reliably, and 2) test the upscaling potential of chambers for EC measurements with varying success in summer- and wintertime (Kade et al., 2012; Maanavilja et al., 2011; Marushchak et al., 2013). In total, EC was used in 11 studies and leaf cuvette measurements 5 (S4).

Based on our findings, we suggest that future studies should incorporate soil respiration measurements into CO<sub>2</sub> flux chamber studies. However, there is no standard way of measuring soil respiration and most of the methods cause disturbance to the ground, or require laboratory work (Kuz'yakov, 2005). Clipping the vegetation was a widely-used method in the investigated articles (Illeris et al., 2004; Nobrega and Grogan, 2008). In some articles, soil respiration was quantified from an non-vegetated soil close to other CO<sub>2</sub> flux measurements (Arens et al., 2008) or with incubation methods (Ylänné et al., 2015). Despite the disturbance caused by clipping, this might be a cost-efficient method for soil respiration measurements, providing there is a sufficient waiting period between the clipping and the measurements (Illeris et al., 2004). Incubation methods will provide robust information on the soil CO<sub>2</sub> emissions for more precise analyses. In addition to soil respiration measurements, the inclusion of non-CO<sub>2</sub> greenhouse gas measurements would provide better understanding of climate forcing, especially in wet regions (CH<sub>4</sub>) and non-vegetated areas (N<sub>2</sub>O) (e.g. Brummell et al., 2012) Furthermore, the incorporation of multiple flux measurement methods will provide more reliable and increasingly detailed information on the spatial and temporal variation of CO<sub>2</sub> fluxes (Myklebust et al., 2008).

**IV Environmental drivers studied to explain fluxes**

GPP, ER and NEE are affected by multiple variables which can be divided into four categories: 1) biota, where the processes occur, 2) resources which are consumed by the biota, 3) regulators which affect the metabolic processes, and 4) disturbance which often destroys the biota and therefore influences the fluxes (Figure 1, 3). Here, we review how these four categories and their variable classes have been included in studies (Table 1). The environmental drivers used to explain fluxes varies depending on the research questions and study designs, yet we assume that with a systematic review (S2) of the environmental drivers measured in the studies, we can highlight overall research gaps.

### *1 Climate*

Air temperature and solar radiation are the most important climatic variables explaining CO<sub>2</sub> fluxes, both spatially and temporally, as temperature controls the metabolic activity of biologic organisms and energy from the sun drives photosynthesis (Lopez-Blanco, et al. 2016, see also Figure 1). Air temperatures and photosynthetically active radiation (PAR, or photon flux density PPFD) in the chamber and/or in the vicinity of loggers are almost always included in the studies, and are often used in the models explaining GPP, ER, and NEE (Karelin et al., 2013; Shaver et al., 2013; Williams et al., 2006).

CO<sub>2</sub> fluxes are driven by soil microclimate which varies spatially over short distances. Soil temperature does not have a direct effect on GPP but it is an important driver of ER (Larsen et al., 2007) as it stimulates soil respiration and decomposition (Grogan and Chapin, 2000). Soil moisture has been identified as a limiting factor for GPP (Nobrega and Grogan, 2008) and for soil and plant respiration, especially at the soil moisture extremes (Dagg and Lafleur, 2011; Giblin et al., 1991; Illeris et al., 2003). The importance of soil microclimate has been recognized in the studies: the temporally varying soil temperature, active layer depth (or thaw depth), soil moisture, and water table height are included in approximately 80 % of the studies (Figure 4). Soil temperature is also often used when modeling ER and NEE, sometimes together with other soil microclimatological variables such as active layer depth or water table height (Christensen et al., 2000; Segal and Sullivan, 2014). Water table height is an important variable, especially in the peatlands, as it controls the anoxic conditions needed for CH<sub>4</sub> emissions.

### *2 Vegetation*

Vegetation plays an important role in controlling the variation of CO<sub>2</sub> fluxes as GPP and, partly, ER are processes occurring within the plant (Johnson et al., 2000). The reviewed

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

364 studies always provide some information on the dominant species with varying degrees of  
365 additional data on vegetation properties (e.g. biomass, LAI). We collected data on the  
366 dominant and secondary species in each measurement plot from the articles and further  
367 divided it into CAVM categories (S2, S5). In many of the studies different measurement  
368 sites within the study area belonged to several CAVM categories.

369  
370 Most of the research has been conducted for the most dominant vegetation types and  
371 species in the Arctic: on graminoid-dominated meadows (with species *Carex* sp.,  
372 *Eriophorum* sp. in >20 of the studies) and prostrate shrub and erect shrub heaths (with  
373 species *Betula nana*, *Empetrum* sp., *Vaccinium vitis-idaea*, *V. uliginosum* in approximately  
374 10-30 of the studies Figure 5A, S5). Fewer studies included cushion plants, forbs, or mosses  
375 and lichens (despite some observations from *Cassiope* sp., *Potentilla* sp., *Sphagnum* sp.,  
376 *Dicranum* sp., *Cladonia* sp., Figure 5B, S5). There are relatively large differences among  
377 measured vegetation types in the most researched areas (Figure 5C). Despite the important  
378 role of *Betula nana*, taller low-shrubs such as *Salix* sp. and in some regions *Juniperus*  
379 *communis* and *Alnus* sp. species are also crucial in the shrubification of the Arctic, yet they  
380 were included in only a small amount of studies. Indeed, based on article keywords, very  
381 few studies aimed to study the effects of shrubification on CO<sub>2</sub> fluxes (Cahoon et al., 2016;  
382 Cahoon et al., 2012). Barren vegetation types, which cover 25 % of the region, are only  
383 included in less than 10 % of the studies, but have received more attention in Antarctica  
384 (Thomazini et al., 2016; Zhu et al., 2014).

385  
386 Although *Eriophorum*, *Carex*, or *Betula nana* -dominated vegetation types have been  
387 relatively well studied they need to be investigated more thoroughly in different  
388 environmental conditions as CO<sub>2</sub> flux observations of tundra shrubs and sedges, together  
389 with moist meadows by streams, contain the largest and most varying CO<sub>2</sub> fluxes (Williams  
390 et al., 2006). Additionally, these are also key sites for CH<sub>4</sub> dynamics. Moreover, we stress  
391 the need for studies focusing on shrubification effects with a diverse combination of taller  
392 low-shrub species. In the CAVM region, low-shrub tundra can be found from European  
393 Russia, northwestern Siberia, and the Yukon-Kuskokwim River delta, as well as in  
394 southwestern parts of Greenland and locally along rivers and tree lines. Regionally, more  
395 observations are needed from wetlands in Alaska and prostrate shrubs in Siberia (4B).  
396 Although having only low CO<sub>2</sub> flux rates, we highlight the need for more studies in barren  
397 soils as these are underrepresented and their spatial coverage might increase resulting  
398 from melting glaciers (Mernild et al., 2012).

399



In addition to plant functional types and plant species, other data on vegetation have been used to understand CO<sub>2</sub> fluxes in different environments (Cornelissen et al., 2003). Plants use carbon for growth and, therefore, plant biomass can be measured to understand carbon sequestration and allocation patterns (mass-ratio hypothesis; Grime, 1998). In addition to biomass measurements, LAI and NDVI have been used to link the amount of photosynthesizing leaves to carbon use patterns, out of which LAI has often been found to be the best predictor of CO<sub>2</sub> dynamics (Marushchak et al., 2013; Shaver et al., 2007; Street et al., 2007). NDVI and LAI measurements were included in approximately 25 studies and biomass sampling in 4 (S4). Plant traits (leaf N, vegetation height, see Dorrepaal 2007) were studied less than biomass (Figure 4), although the relationship between leaf N and CO<sub>2</sub> fluxes has received increasing attention (Street et al., 2012).

Based on our findings, we firstly highlight the need for more detailed analysis of vegetation properties and CO<sub>2</sub> fluxes. For example, the relationship between plant traits or growth and CO<sub>2</sub> fluxes remains uncertain (De Deyn et al., 2008). Additionally, there is a need to study the linkages of ER and biomass, as higher biomass is linked to higher GPP but not as clearly as to ER (Boelman et al., 2003; Douma et al., 2007; Poyatos et al., 2014). Lastly, we stress the need for vegetation data -based upscaling, particularly to landscape scales, which remains a challenge in carbon cycle studies (O'Rourke et al., 2015). Although chamber-derived GPP, ER, and NEE have been linked to field-observations of spectral properties of vegetation and LAI (Boelman et al., 2003; Williams et al., 2006), this relationship has seldom been used to upscale results to larger areas with aerial or satellite imagery (Marushchak et al., 2013). More research of CO<sub>2</sub> fluxes and remote sensing -based indices at multiple resolutions (e.g. WorldView, Landsat, MODIS, see e.g. Curasi et al., 2016) is called for to develop the upscaling methodology. Recent progress of unmanned aerial vehicles (UAVs) offers CO<sub>2</sub> flux researchers new opportunities for fine spatial and hyperspectral resolution imagery data (Anderson and Gaston, 2013). To our knowledge, there are no published studies utilizing UAVs for CO<sub>2</sub> flux chamber measurements in the tundra.

### *3 Soils*

GPP, ER and NEE are regulated by soil physical, chemical, and biological properties which vary greatly over short distances (Arnesen et al., 2007; Wang et al., 2010). Soil physical properties, particularly soil type, texture, and organic layer depth influence overall soil conditions (Buchmann, 2000; Grand et al., 2016) and are connected to soil temperature, depth of the active layer, and vegetation properties. Soil nutrients are an important

1  
2  
3 437 resource for plants and soil microbes, therefore affecting GPP and ER (Grogan and  
4 438 Jonasson, 2005). Soil N together with P limit tundra productivity, leading to increases in  
5 439 GPP if nutrients are added to the soil (Arens et al., 2008; Shaver et al., 1998). Soil microbes  
6 440 (heterotrophic respiration) control soil respiration together with plant roots (autotrophic  
7 441 respiration) and, therefore, partly ER (Segal and Sullivan, 2014). Soil data is included in  
8 442 fewer studies than climate and vegetation data: soil type descriptions are included in 15  
9 443 studies, litter biomass estimates in 7 studies, soil nitrogen data in 18, and SOC  
10 444 measurements in 15 of the studies (S4). Data from the soil microbes is included in eight  
11 445 studies.  
12  
13  
14  
15  
16  
17

18 447 We recommend future studies to at least describe the soil type (Jones et al., 2010) and  
19 448 organic layer depth to provide information on soil physical and chemical conditions (see  
20 449 e.g. Biasi et al., 2008). The inclusion of soil data is needed in spatial study designs as soil  
21 450 properties vary especially spatially. We highlight the need for better understanding of CO<sub>2</sub>  
22 451 flux – nutrient and SOC relationships (Williams et al., 2006) as recent studies have  
23 452 discovered the respiratory losses and carbon sequestration to soils to depend on soil  
24 453 geochemistry (Doetterl et al., 2015). Thus, including soil nutrient data, of for example N  
25 454 and P, also increases knowledge of the linkages between different biogeochemical cycles  
26 455 in the Arctic. We stress the need for easily measurable indicators of soil quality (e.g. soil  
27 456 pH, the color or depth of organic matter, e.g. White et al., 2004). Microbial activity should  
28 457 receive more attention as it is not yet fully understood how microbial communities and  
29 458 their functioning regulate ER and NEE and how they will respond to environmental change  
30 459 (Biasi et al., 2008; Hultman et al., 2015; Xue et al., 2016).  
31  
32  
33  
34  
35  
36  
37  
38

39 461 *4 Disturbance*  
40  
41 462

42  
43 463 Geomorphological processes create disturbance that leads to heterogeneous abiotic and  
44 464 biotic conditions (le Roux and Luoto, 2014) potentially shaping CO<sub>2</sub> fluxes (Vogel et al.,  
45 465 2009). There is a small but increasing amount of publications that focus on the effect of  
46 466 fluvial processes (Curasi et al., 2016; Oberbauer et al., 2007), cryoturbation as in e.g.  
47 467 patterned ground and hummocks (Heikkinen et al., 2002; Maanavilja et al., 2011), and  
48 468 permafrost as in e.g. palsa mires (Bäckstrand et al., 2010; Nykanen et al., 2003) on CO<sub>2</sub>  
49 469 fluxes. Most of these studies have been conducted in northern Fennoscandia, though there  
50 470 is an increasing number of studies of permafrost thaw from Alaska (Natali et al., 2011;  
51 471 Schuur et al., 2009).  
52  
53  
54  
55  
56  
57  
58  
59  
60



CO<sub>2</sub> fluxes are mediated by biological disturbance (Falk et al., 2015; Sjögersten et al., 2011; Susiluoto et al., 2008). The tundra is grazed by large and small herbivores which can alter species composition, above- and belowground biomass and litter quality and, therefore, CO<sub>2</sub> fluxes (Cahoon et al., 2012; Metcalfe and Olofsson, 2015; Ylanne et al., 2015). Herbivory was included in 13 studies with a focus on larger herbivores such as reindeer and muskoxen (S4). Larval outbreak was included in one study (but see a new study by Lund et al., 2017). In addition to herbivory, disease outbreaks (plant pathogens) can modify CO<sub>2</sub> fluxes but they have only been studied to a rather limited extent in one study. Olofsson et al., (2011) demonstrated GPP to decrease during the growing season due to an outbreak of a parasitic fungus, *Arwidssonia empetri* in northern Sweden. Presently, intensive insect outbreaks are rare in the tundra, but this might change if the climate warms (Abbot et al., 2016).

Fire is an important disturbance in boreal regions, but has had only minor effects on tundra carbon cycling since the early Holocene (Kasischke and Turetsky, 2006). However, models predict an increasing amount of tundra fires in a warmer climate which has already been observed in Alaska (Hu et al., 2010; Krawchuk et al., 2009), but still the effect of fire disturbance on tundra carbon cycling has received relatively little attention. Although there are studies considering the SOC stocks before and after a fire (Bret-Harte et al., 2013; Mack et al., 2011), there are no CO<sub>2</sub> flux chamber studies related to fire in the tundra (but see e.g. Köster et al., 2016 on boreal regions).

The disturbance regime of the studied region is often well described in the study site description (e.g. Vourtilis et al., 2000), but it should be better incorporated into analyses due to predicted changes in occurrence in a warmer climate (Aalto et al., 2014; Koven et al., 2011). Moreover, we need more comprehensive analysis of the effect of disturbance in different conditions (e.g. in nutrient-rich vs. nutrient-poor environments, or high vs. low-Arctic) as the response of an ecosystem to e.g. drought, increased herbivory or fire differs depending on the location.

## V Towards a conceptual framework

Conceptual frameworks to explain and describe the drivers and mechanisms of fine-scale spatio-temporal variation of CO<sub>2</sub> fluxes are lacking (Bardgett and Wardle, 2010; Callaghan et al., 2004; Xue et al., 2016). Tundra ecosystems are complex and heterogeneous, so to be able to predict future changes, we first need to create a good understanding on the current processes and variables affecting CO<sub>2</sub> fluxes. Firstly, we highlight the need for

theoretical, analytical, and process-based justification for the choice of measured variables. Secondly, we stress the importance of understanding the direct and indirect relationships among the variables (for example, the importance of topography, Figure 3). Most of the reviewed publications include data on biota (vegetation), resources (radiation, soil moisture), and regulators (air and/or soil temperature) as these variables vary greatly both spatially and temporally (S4). Out of these drivers, air temperature, PAR, vegetation type, LAI and, occasionally, soil temperature and moisture and NDVI were the most important factors affecting CO<sub>2</sub> fluxes. Other environmental variables measured depend on the study location and design, and research questions. Measuring disturbance is challenging as disturbance often occurs as episodic events. Still, we encourage the investigators to describe the disturbance regime of the plot by for example collecting herbivory data or describing the active geomorphological processes.

*1 Standardized methodology for site description of chamber studies*

The description of the measured sites varies between the studies. Both species-level as well as plant functional type (PFT) descriptions are used in the literature. For example, classifications of meadow-heath (Oberbauer et al., 2007) and more precise groups based on the dominant species i.e. *Betula nana* and *Dryas octopetala* (Shaver et al., 2007) have been used in the studies. Because of the importance of vegetation in the carbon cycle, standardized and easily comparable descriptive information on vegetation is crucial in order to understand CO<sub>2</sub> fluxes. Although the vegetation data given by CO<sub>2</sub> flux researchers is descriptive, we encourage the investigators to use widely used plant community classifications in addition to their own. A comprehensive system for the grouping of plants with similar links to environmental conditions can reduce the complexity of the systems (Otieno et al., 2009). Here, we used CAVM (Walker et al., 2005) which was developed for biogeographic and landscape-scale purposes. It has been successfully applied to the carbon cycle studies and allows researchers to easily compare results (Arndal et al., 2009; Biasi et al., 2008; Douma et al., 2007). An even more precise hierarchical floristic classification system than CAVM which is based on a smaller habitat-scale has been developed by Mucina et al. (2016), but is currently available only for Europe. In addition, the Arctic Vegetation Archive (AVA) aims to combine Arctic vegetation data at plot-scale but is still under progress (Walker et al., 2016).

Despite a common grouping system and characteristic vegetation types, smaller wet patches, stream channels and transitional zones are equally important in the patchy tundra vegetation. Fletcher et al. (2012) have shown that the most abundant vegetation types (e.g.

*Betula nana* or *Empetrum* heath) have a higher GPP than their mixed vegetation types (e.g. *Betula-Empetrum* heath). Therefore, measuring also less abundant transition zones and providing species cover data on the dominant and secondary species to locate them is important. Currently, species cover data is included in approximately only 50 % of the studies, and cryptogram (mosses and lichens) data in even less. Cryptograms cover large areas in the Arctic, and the crucial role of mosses in the carbon cycle has been identified, especially during spring when leaves of deciduous vascular plants are not yet fully developed but mosses are already photosynthesizing (Douma et al., 2007). Cryptogram quantification using remote sensing, and LAI determination, are much less studied as they are methodologically even more challenging than vascular plants. However, high-resolution satellite and UAV imagery might provide tools for robust vegetation characterization in the future (Fraser et al., 2016). In addition to standardized vegetation characterization, chamber researchers need a common platform, similar to Fluxnet created for EC studies (Baldocchi et al., 2001; Chu et al., 2017), to share data and develop methods and guidelines.

## VI Summary and future research needs

Environmental variation can affect tundra CO<sub>2</sub> fluxes in multiple ways. These effects are not yet fully understood due to limitations posed by study locations, designs and environmental variables measured to explain chamber-derived CO<sub>2</sub> fluxes (Table 2). Even though study locations cover the environmental variation relatively well, more research is needed in extreme conditions (such as high precipitation and low temperatures, high topsoil pH, high SOC stock conditions). Study designs should focus on covering spatial variation in environmental conditions, vegetation and soils, within and between study areas. Additionally, both ends of the productivity gradient (low-shrubs and barren vegetation types) should be studied more thoroughly.

Ecologically and physically relevant variables have been included in the reviewed studies to varying degrees and some topics require new research more urgently than others. Chamber-derived CO<sub>2</sub> fluxes should be explained not only by microclimate and vegetation type, but also by other fluxes, soil microbes and nutrient availability, and disturbance data, to better understand ecosystem functioning. As tundra regions are typically nutrient-limited thus responding strongly to changes in nutrient availability, more studies on the relationship of soil nutrients and microbes with GPP, ER, and NEE measurements are urgently needed. Simultaneous measurements of other greenhouse gases and soil respiration together with CO<sub>2</sub> fluxes are particularly urgent themes as changes therein might be crucial in determining the future carbon balance.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

584 Finally, studies of CO<sub>2</sub> fluxes should focus on the sensitive processes – along with their  
585 transitions and thresholds in the ecosystem – that have the potential to change the carbon  
586 balance. According to an expert assessment of the most important processes in the  
587 changing Arctic environment (Abbott, et al. 2016), these sensitive processes include  
588 vegetation shifts, permafrost thaw as well as hydrological changes. The effect of  
589 permafrost thaw has received increasing attention, but it is still unclear how other  
590 geomorphological disturbance, fire events, shrubification, or disease and herbivory  
591 outbreaks influence CO<sub>2</sub> fluxes and the tundra carbon balance. It is thus crucial to deepen  
592 the understanding of these processes. A better conceptual framework and understanding  
593 of CO<sub>2</sub> fluxes will provide more information on the carbon cycling in the tundra.

For Peer Review

## References

- Aalto J, Le Roux PC and Luoto M (2013) Vegetation Mediates Soil Temperature and Moisture in Arctic-Alpine Environments. *Source: Arctic, Antarctic, and Alpine Research Published By: Institute of Arctic and Alpine Research Arctic, Antarctic, and Alpine Research* 45(4): 429–439.
- Aalto J, Venäläinen A, Heikkinen RK, et al. (2014) Potential for extreme loss in high-latitude Earth surface processes due to climate change. *Geophysical Research Letters* 41(11): 3914–3924.
- Abbott BW, Jones JB, G Schuur EA, et al. (2016) Biomass offsets little or none of permafrost carbon release from soils, streams, and wildfire: an expert assessment. *Environmental Research Letters* 11(3): 1–13.
- Anderson K and Gaston KJ (2013) Lightweight unmanned aerial vehicles will revolutionize spatial ecology. *Frontiers in Ecology and the Environment* 11(3): 138–146.
- Arens SJT, Sullivan PF and Welker JM (2008) Nonlinear responses to nitrogen and strong interactions with nitrogen and phosphorus additions drastically alter the structure and function of a high arctic ecosystem. *Journal of Geophysical Research-Biogeosciences* 113(G3): 10.
- Arndal MF, Illeris L, Michelsen A, et al. (2009) Seasonal Variation in Gross Ecosystem Production, Plant Biomass, and Carbon and Nitrogen Pools in Five High Arctic Vegetation Types. *Arctic Antarctic and Alpine Research* 41(2): 164–173.
- Arnesen G, Bech PSA and Engelskjøn T (2007) Soil acidity, content of carbonates, and available phosphorus are the soil factors best correlated with alpine vegetation: evidence from Troms, north Norway. *Arctic, Antarctic, and Alpine Research* 39(2): 189–199.
- Asner GP, Scurlock JMO and A. Hicke J (2003) Global synthesis of leaf area index observations: implications for ecological and remote sensing studies. *Global Ecology and Biogeography* 12(3): 191–205.
- Baldocchi D, Falge E, Gu L, et al. (2001) FLUXNET: A New Tool to Study the Temporal and Spatial Variability of Ecosystem-Scale Carbon Dioxide, Water Vapor, and Energy Flux Densities. *Bulletin of the American Meteorological Society* 82(11): 2415–2434.
- Bardgett RD and Wardle DA (2010) *Aboveground-belowground linkages: biotic interactions, ecosystem processes, and global change*. Oxford University Press.
- Belshe EF, Schuur EAG and Bolker BM (2013) Tundra ecosystems observed to be CO<sub>2</sub> sources due to differential amplification of the carbon cycle. *Ecology Letters* 16(10): 1307–1315.
- Biasi C, Meyer H, Rusalimova O, et al. (2008) Initial effects of experimental warming on carbon exchange rates, plant growth and microbial dynamics of a lichen-rich dwarf shrub tundra in Siberia. *Plant and Soil* 307(1–2): 191–205.
- Biasi Christina, Meyer H, Rusalimova O, et al. (2008) Initial effects of experimental warming on carbon exchange rates, plant growth and microbial dynamics of a lichen-rich dwarf shrub tundra in Siberia. *Plant and Soil* 307(1–2): 191–205.
- Billings WD (1973) Arctic and Alpine Vegetations: Similarities, Differences, and Susceptibility to Disturbance. *BioScience* 23(12): 697–704.

1  
2  
3 651 Billings WD and Mooney HA (1968) The ecology of arctic and alpine plants. *Biological*  
4 652 *Reviews* 43(4): 481–529.  
5 653 Boelman NT, Stieglitz M, Rueth HM, et al. (2003) Response of NDVI, biomass, and  
6 654 ecosystem gas exchange to long-term warming and fertilization in wet sedge tundra.  
7 655 *Oecologia* 135(3): 414–421.  
8 656 Bokhorst S, Bjerke JW, Melillo J, et al. (2010) Impacts of extreme winter warming events on  
9 657 litter decomposition in a sub-Arctic heathland. *Soil Biology and Biochemistry* 42(4):  
10 658 611–617.  
11 659 Bret-Harte MS, Mack MC, Shaver GR, et al. (2013) The response of Arctic vegetation and  
12 660 soils following an unusually severe tundra fire. *Philosophical Transactions of the Royal*  
13 661 *Society of London B: Biological Sciences* 368(1624): 1–15.  
14 662 Brummell ME, Farrell RE and Siciliano SD (2012) Greenhouse gas soil production and  
15 663 surface fluxes at a high arctic polar oasis. *Soil Biology and Biochemistry* 52: 1–12.  
16 664 Buchmann N (2000) Biotic and abiotic factors controlling soil respiration rates in Picea  
17 665 abies stands. *Soil Biology & Biochemistry* 32(11–12): 1625–1635.  
18 666 Bäckstrand K, Crill PM, Mastepanov M, et al. (2008) Non-methane volatile organic  
19 667 compound flux from a subarctic mire in Northern Sweden. *Tellus B: Chemical and*  
20 668 *Physical Meteorology* 60(2): 226–237.  
21 669 Bäckstrand K, Crill PM, Jackowicz-Korcz Nski M, et al. (2010) Annual carbon gas budget for  
22 670 a subarctic peatland, Northern Sweden. *Biogeosciences* 7(1): 95–108.  
23 671 Cahoon S. M. P., Sullivan PF, Shaver GR, et al. (2012) Interactions among shrub cover and  
24 672 the soil microclimate may determine future Arctic carbon budgets. *Ecology Letters*  
25 673 15(12): 1415–1422.  
26 674 Cahoon Sean M P, Sullivan PF, Shaver GR, et al. (2012) Interactions among shrub cover and  
27 675 the soil microclimate may determine future Arctic carbon budgets. *Ecology Letters* 15:  
28 676 1415–1422.  
29 677 Cahoon S.M.P., Sullivan PF, Post E, et al. (2012) Large herbivores limit CO<sub>2</sub> uptake and  
30 678 suppress carbon cycle responses to warming in West Greenland. *Global Change*  
31 679 *Biology* 18(2): 469–479.  
32 680 Cahoon SP, Sullivan P and Post E (2016) Greater Abundance of Betula nana and Early Onset  
33 681 of the Growing Season Increase Ecosystem CO<sub>2</sub> Uptake in West Greenland. *Ecosystems*.  
34 682 Callaghan T V, Björn LO, Chernov Y, et al. (2004) Effects on the Function of Arctic  
35 683 Ecosystems in the Short-and Long-term Perspectives Climate Change and UV-B  
36 684 Impacts on Arctic Tundra and Polar Desert Ecosystems Effects on the Function of  
37 685 Arctic Ecosystems in the Short-and Long-term Perspectives. *AMBIO* 33(7): 448–458.  
38 686 Christensen TR, Friborg T, Sommerkorn M, et al. (2000) Trace gas exchange in a high-arctic  
39 687 valley 1. Variations in CO<sub>2</sub> and CH<sub>4</sub> flux between tundra vegetation types. *Global*  
40 688 *Biogeochemical Cycles* 14(3): 701–713.  
41 689 Chu H, Baldocchi DD, John R, et al. (2017) Fluxes All of the Time? A Primer on the Temporal  
42 690 Representativeness of FLUXNET. *Journal of Geophysical Research: Biogeosciences*  
43 691 122(2): 289–307.  
44 692 Cornelissen JHC, Lavorel S, Garnier E, et al. (2003) A handbook of protocols for  
45 693 standardised and easy measurement of plant functional traits worldwide. *Australian*



- 694 *Journal of Botany* 51(4): 335–380.
- 695 Curasi SR, Loranty MM and Natali SM (2016) Water track distribution and effects on carbon  
696 dioxide flux in an eastern Siberian upland tundra landscape. *Environmental Research*  
697 *Letters* 11(4): 45002.
- 698 Dagg J and Lafleur P (2010) An application of plot-scale NDVI in predicting carbon dioxide  
699 exchange and leaf area index in heterogeneous subarctic tundra. *Canadian Journal of*  
700 *Remote Sensing* 36: S111–S123.
- 701 Dagg J and Lafleur P (2011) Vegetation Community, Foliar Nitrogen, and Temperature  
702 Effects on Tundra CO<sub>2</sub> Exchange across a Soil Moisture Gradient. *Arctic Antarctic and*  
703 *Alpine Research* 43(2): 189–197.
- 704 De Deyn GB, Cornelissen JHC and Bardgett RD (2008) Plant functional traits and soil carbon  
705 sequestration in contrasting biomes. *Ecology Letters* 11(5): 516–531.
- 706 Dinerstein E, Olson D, Joshi A, et al. (2017) An Ecoregion-Based Approach to Protecting  
707 Half the Terrestrial Realm. *BioScience* 67(6): 534–545.
- 708 Doetterl S, Stevens A, Six J, et al. (2015) Soil carbon storage controlled by interactions  
709 between geochemistry and climate. *Nature Geoscience, Nature Research* 8(10): 780–  
710 783.
- 711 Dorrepaal E (2007) Are plant growth-form-based classifications useful in predicting  
712 northern ecosystem carbon cycling feedbacks to climate change? *Journal of Ecology*  
713 95(6): 1167–1180.
- 714 Douma JC, Van Wijk MT, Lang SI, et al. (2007) The contribution of mosses to the carbon  
715 and water exchange of arctic ecosystems: quantification and relationships with system  
716 properties. *Plant Cell and Environment* 30(10): 1205–1215.
- 717 Elmendorf SC, Henry GHR, Hollister RD, et al. (2012) Plot-scale evidence of tundra  
718 vegetation change and links to recent summer warming. *Nature Climate Change* 2(6):  
719 453–457.
- 720 Falk JM, Schmidt NM, Christensen TR, et al. (2015) Large herbivore grazing affects the  
721 vegetation structure and greenhouse gas balance in a high arctic mire. *Environmental*  
722 *Research Letters* 10(4): 11.
- 723 Flessa H, Rodionov A, Guggenberger G, et al. (2008) Landscape controls of CH<sub>4</sub> fluxes in a  
724 catchment of the forest tundra ecotone in northern Siberia. *Global Change Biology*  
725 14(9): 1–17.
- 726 Fletcher BJ, Gornall JL, Poyatos R, et al. (2012) Photosynthesis and productivity in  
727 heterogeneous arctic tundra: Consequences for ecosystem function of mixing  
728 vegetation types at stand edges. *Journal of Ecology* 100(2): 441–451.
- 729 Forbes BC, Fauria MM and Zetterberg P (2010) Russian Arctic warming and ‘greening’ are  
730 closely tracked by tundra shrub willows. *Global Change Biology* 16(5): 1542–1554.
- 731 Fraser RH, Olthof I, Lantz TC, et al. (2016) UAV photogrammetry for mapping vegetation in  
732 the low-Arctic. *Arctic Science, NRC Research Press* <http://www.nrcresearchpress.com>  
733 2(3): 79–102.
- 734 Giblin AE, Nadelhoffer KJ, Shaver GR, et al. (1991) Biogeochemical Diversity Along a  
735 Riverside Toposequence in Arctic Alaska. *Ecological Monographs* 61(4): 415–435.
- 736 Gil J, Pérez T, Boering K, et al. (2017) Mechanisms responsible for high N<sub>2</sub>O emissions from



1  
2  
3 737 subarctic permafrost peatlands studied via stable isotope techniques. *Global*  
4 738 *Biogeochemical Cycles* 31(1): 172–189.  
5 739 Grand S, Rubin A, Verrecchia EP, et al. (2016) Variation in Soil Respiration across Soil and  
6 740 Vegetation Types in an Alpine Valley. *PLOS ONE* 11(9): 1–16.  
7 741 Griffis TJ, Rouse WR and Waddington JM (2000) Scaling net ecosystem CO<sub>2</sub> exchange from  
8 742 the community to landscape-level at a subarctic fen. *Global Change Biology* 6(4): 459–  
9 743 473.  
10 744 Grime JP (1998) Benefits of plant diversity to ecosystems: immediate, filter and founder  
11 745 effects. *Journal of Ecology* 86(6): 902–910.  
12 746 Grogan P and Chapin FS (2000) Initial effects of experimental warming on above- and  
13 747 belowground components of net ecosystem CO<sub>2</sub> exchange in arctic tundra. *Oecologia*  
14 748 125(4): 512–520.  
15 749 Grogan P and Jonasson S (2005) Temperature and substrate controls on intra-annual  
16 750 variation in ecosystem respiration in two subarctic vegetation types. *Global Change*  
17 751 *Biology* 11(3): 465–475.  
18 752 Hartley IP, Garnett MH, Sommerkorn M, et al. (2012) A potential loss of carbon associated  
19 753 with greater plant growth in the European Arctic. *Nature Climate Change* 2: 875–870.  
20 754 Heikkinen JEP, Maljanen M, Aurela M, et al. (2002) Carbon dioxide and methane dynamics  
21 755 in a sub-Arctic peatland in northern Finland. *Polar Research* 21(1): 49–62.  
22 756 Hinzman LD, Bettez ND, Bolton WR, et al. (2005) Evidence and Implications of Recent  
23 757 Climate Change in Northern Alaska and Other Arctic Regions. *Climatic Change* 72(3):  
24 758 251–298.  
25 759 Hinzman LD, Deal CJ, McGuire AD, et al. (2013) Trajectory of the Arctic as an integrated  
26 760 system. *Ecological Applications* 23(8): 1837–1868.  
27 761 Hobbie SE (1996) Temperature and plant species control over litter decomposition in  
28 762 Alaskan tundra. *Ecological Monographs* 66(664): 503–522.  
29 763 Hu FS, Higuera PE, Walsh JE, et al. (2010) Tundra burning in Alaska: Linkages to climatic  
30 764 change and sea ice retreat. *Journal of Geophysical Research* 115(G4): 1–8.  
31 765 Hudson JMG and Henry GHR (2009) Increased plant biomass in a high arctic heath  
32 766 community from 1981 to 2008. *Ecology* 90(10): 2657–2663.  
33 767 Hugelius G, Strauss J, Zubrzycki S, et al. (2014) Estimated stocks of circumpolar permafrost  
34 768 carbon with quantified uncertainty ranges and identified data gaps. *Biogeosciences*  
35 769 11(23): 6573–6593.  
36 770 Hultman J, Waldrop MP, Mackelprang R, et al. (2015) Multi-omics of permafrost, active  
37 771 layer and thermokarst bog soil microbiomes. *Nature* 521(7551): 208–212.  
38 772 Hutchinson GL and Livingston GP (2001) Vents and seals in non-steady-state chambers  
39 773 used for measuring gas exchange between soil and the atmosphere. *European Journal*  
40 774 *of Soil Science* 52(4): 675–682.  
41 775 Illeris L, Michelsen A and Jonasson S (2003) Soil plus root respiration and microbial  
42 776 biomass following water, nitrogen, and phosphorus application at a high arctic semi  
43 777 desert. *Biogeochemistry* 65(1): 15–29.  
44 778 Illeris L, Konig SM, Grogan P, et al. (2004) Growing-season carbon dioxide flux in a dry  
45 779 subarctic heath: Responses to long-term manipulations. *Arctic Antarctic and Alpine*

- 780 *Research* 36(4): 456–463.
- 781 Johnson LC, Shaver GR, Cades DH, et al. (2000) Plant carbon-nutrient interactions control  
782 CO<sub>2</sub> exchange in Alaskan wet sedge tundra ecosystems. *Ecology* 81(2): 453–469.
- 783 Kade A, Bret-Harte MS, Euskirchen ES, et al. (2012) Upscaling of CO<sub>2</sub> fluxes from  
784 heterogeneous tundra plant communities in Arctic Alaska. *Journal of Geophysical*  
785 *Research - Biogeosciences* 117(G4): G04007.
- 786 Karelin D V., Zamolodchikov DG, Zukert N V., et al. (2013) Interannual changes in PAR and  
787 soil moisture during the warm season may be more important for directing of annual  
788 carbon balance in tundra than temperature fluctuations. *Zhurnal Obshchei Biologii*  
789 74(1): 3–22.
- 790 Kasischke ES and Turetsky MR (2006) Recent changes in the fire regime across the North  
791 American boreal region—Spatial and temporal patterns of burning across Canada and  
792 Alaska. *Geophysical Research Letters* 33(9): 1–5.
- 793 Kelsey KC, Leffler AJ, Beard KH, et al. (2016) Interactions among climate, topography and  
794 herbivory control greenhouse gas (CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O) fluxes in a subarctic coastal  
795 wetland. *Journal of Geophysical Research - Biogeosciences* 121(12): 2960–2975.
- 796 Knowles JF, Blanken PD and Williams MW (2016) Wet meadow ecosystems contribute the  
797 majority of overwinter soil respiration from snow-scoured alpine tundra. *Journal of*  
798 *Geophysical Research: Biogeosciences* 121: 1118–1130.
- 799 Koven CD, Ringeval B, Friedlingstein P, et al. (2011) Permafrost carbon-climate feedbacks  
800 accelerate global warming. *PNAS* 108(36): 14769–14774.
- 801 Krawchuk MA, Moritz MA, Parisien M-A, et al. (2009) Global Pyrogeography: the Current  
802 and Future Distribution of Wildfire. Chave J (ed.), *PLoS ONE* 4(4): 1–12.
- 803 Kuhry P, Dorrepaal E, Hugelius G, et al. (2010) Short communication: Potential  
804 remobilization of belowground permafrost carbon under future global warming.  
805 *Permafrost and Periglacial Processes* 21(2): 208–214.
- 806 Kutzbach L, Schneider J, Sachs T, et al. (2007) CO<sub>2</sub> flux determination by closed-chamber  
807 methods can be seriously biased by inappropriate application of linear regression.  
808 *Biogeosciences* 4: 1005–1025.
- 809 Kuzyakov Y (2005) Sources of CO<sub>2</sub> efflux from soil and review of partitioning methods.
- 810 Köster K, Köster E, Orumaa A, et al. (2016) How time since forest fire affects stand structure,  
811 soil physical-chemical properties and soil CO<sub>2</sub> efflux in hemiboreal scots pine forest  
812 fire chronosequence? *Forests* 7(9): 1–12.
- 813 Larsen KS, Ibrom A, Jonasson S, et al. (2007) Significance of cold-season respiration and  
814 photosynthesis in a subarctic heath ecosystem in Northern Sweden. *Global Change*  
815 *Biology* 13(7): 1498–1508.
- 816 le Roux PC and Luoto M (2014) Earth surface processes drive the richness, composition and  
817 occurrence of plant species in an arctic-alpine environment. *Journal of Vegetation*  
818 *Science* 25(1): 45–54.
- 819 Leffler AJ, Klein ES, Oberbauer SF, et al. (2016) Coupled long-term summer warming and  
820 deeper snow alters species composition and stimulates gross primary productivity in  
821 tussock tundra. *Oecologia* 181(1): 287–297.
- 822 López-Blanco E, Lund M, Williams M, et al. (2016) Exchange of CO<sub>2</sub> in Arctic tundra:

1  
2  
3 823 impacts of meteorological variations and biological disturbance. *Biogeosciences*  
4 824 *Discussions* doi:10.519.  
5 825 Lund M, Raundrup K, Westergaard-Nielsen A, et al. (2017) Larval outbreaks in West  
6 826 Greenland: Instant and subsequent effects on tundra ecosystem productivity and CO<sub>2</sub>  
7 827 exchange. *Ambio* 46: 26–38.  
8 828 Luo Y, Ahlström A, Allison SD, et al. (2016) Toward more realistic projections of soil carbon  
9 829 dynamics by Earth system models. *Global Biogeochemical Cycles* 30(1): 40–56.  
10 830 Maanavilja L, Riutta T, Aurela M, et al. (2011) Spatial variation in CO<sub>2</sub> exchange at a  
11 831 northern aapa mire. *Biogeochemistry* 104(1–3): 325–345.  
12 832 Mack MC, Sydoniabret-Harte M, Hollingsworth T, et al. (2011) Carbon loss from an  
13 833 unprecedented Arctic tundra wildfire. *Nature* 475(7357): 489–492.  
14 834 Marushchak ME, Pitkämäki A, Koponen H, et al. (2011) Hot spots for nitrous oxide  
15 835 emissions found in different types of permafrost peatlands. *Global Change Biology*  
16 836 17(8): 2601–2614.  
17 837 Marushchak ME, Kiepe I, Biasi C, et al. (2013) Carbon dioxide balance of subarctic tundra  
18 838 from plot to regional scales. *Biogeosciences* 10(1): 437–452.  
19 839 McGuire AD, Anderson LG, Christensen TR, et al. (2009) Sensitivity of the carbon cycle in  
20 840 the Arctic to climate change. *Ecological Monographs* 79(4): 523–555.  
21 841 McGuire A. D., Hayes DJ, Kicklighter DW, et al. (2010b) An analysis of the carbon balance of  
22 842 the Arctic Basin from 1997 to 2006. *Tellus, Series B: Chemical and Physical Meteorology*.  
23 843 McGuire A. David, Macdonald RW, Schuur EAG, et al. (2010a) The carbon budget of the  
24 844 northern cryosphere region. *Current Opinion in Environmental Sustainability*.  
25 845 McGuire AD, Christensen TR, Hayes D, et al. (2012) An assessment of the carbon balance of  
26 846 Arctic tundra: comparisons among observations, process models, and atmospheric  
27 847 inversions. *Biogeosciences* 9(8): 3185–3204.  
28 848 Mernild SH, Liston GE, Mernild SH, et al. (2012) Greenland Freshwater Runoff. Part II:  
29 849 Distribution and Trends, 1960–2010. *Journal of Climate* 25(17): 6015–6035.  
30 850 Metcalfe DB and Olofsson J (2015) Distinct impacts of different mammalian herbivore  
31 851 assemblages on arctic tundra CO<sub>2</sub> exchange during the peak of the growing season.  
32 852 *Oikos* 124(12): 1632–1638.  
33 853 Mucina L, Bültmann H, Dierßen K, et al. (2016) Vegetation of Europe: hierarchical floristic  
34 854 classification system of vascular plant, bryophyte, lichen, and algal communities.  
35 855 *Applied Vegetation Science* 19(S1): 3–264.  
36 856 Myklebust MC, Hipps LE and Ryel RJ (2008) Comparison of eddy covariance, chamber, and  
37 857 gradient methods of measuring soil CO<sub>2</sub> efflux in an annual semi-arid grass, *Bromus*  
38 858 *tectorum*. *Agricultural and Forest Meteorology* 148(11): 1894–1907.  
39 859 Nadelhoffer KJ, Giblin AE, Shaver GR, et al. (1991) Effects of Temperature and Substrate  
40 860 Quality on Element Mineralization in Six Arctic Soils. *Ecology* 72(1): 242–253.  
41 861 Natali SM, Schuur EAG, Trucco C, et al. (2011) Effects of experimental warming of air, soil  
42 862 and permafrost on carbon balance in Alaskan tundra. *Global Change Biology* 17(3):  
43 863 1394–1407.  
44 864 Nobrega S and Grogan P (2008) Landscape and ecosystem-level controls on net carbon  
45 865 dioxide exchange along a natural moisture gradient in Canadian low arctic tundra.

- 866 *Ecosystems* 11(3): 377–396.
- 867 Nykanen H, Heikkinen JEP, Pirinen L, et al. (2003) Annual CO<sub>2</sub> exchange and CH<sub>4</sub> fluxes on a  
868 subarctic tundra mire during climatically different years. *Global Biogeochemical Cycles*  
869 17(1): 18.
- 870 O'Rourke SM, Angers DA, Holden NM, et al. (2015) Soil organic carbon across scales.  
871 *Global Change Biology* 21(10): 3561–3574.
- 872 Oberbauer SF, Tweedie CE, Welker JM, et al. (2007) Tundra CO<sub>2</sub> fluxes in response to  
873 experimental warming across latitudinal and moisture gradients. *Ecological*  
874 *Monographs* 77(2): 221–238.
- 875 Oechel WC, Hastings SJ, Vourlitis G, et al. (1993) Recent change of Arctic tundra  
876 ecosystems from a net carbon dioxide sink to a source. *Nature* 361(6412): 520–523.
- 877 Oechel WC, Vourlitis G L., Hastings SJ, et al. (2000) Acclimation of ecosystem CO<sub>2</sub> exchange  
878 in the Alaskan Arctic in response to decadal climate warming. *Nature* 406(6799): 978–  
879 81.
- 880 Olivas PC, Oberbauer SF, Tweedie CE, et al. (2010) Responses of CO<sub>2</sub> flux components of  
881 Alaskan Coastal Plain tundra to shifts in water table. *Journal of Geophysical Research-*  
882 *Biogeosciences* 115(G4): 1–13.
- 883 Olofsson J, Ericson L, Torp M, et al. (2011) Carbon balance of Arctic tundra under increased  
884 snow cover mediated by a plant pathogen. *Nature Climate Change* 1(4): 220–223.
- 885 Olson DM, Dinerstein E, Wikramanayake ED, et al. (2001) Full-Text. *BioScience* 51(11).
- 886 Osterkamp TE and Romanovsky VE (1999) Evidence for warming and thawing of  
887 discontinuous permafrost in Alaska. *Permafrost and Periglacial Processes* 10(1): 17–37.
- 888 Otieno DO, Wartinger M, Nishiwaki A, et al. (2009) Responses of CO<sub>2</sub> Exchange and Primary  
889 Production of the Ecosystem Components to Environmental Changes in a Mountain  
890 Peatland. *Ecosystems* 12(4): 590–603.
- 891 Overeem I and Syvitski JPM (2010) Shifting Discharge Peaks in Arctic Rivers, 1977–2007.  
892 *Geografiska Annaler Series A Physical Geography* 92(2): 285–296.
- 893 Pare MC and Bedard-Haughn A (2012) Landscape-scale N mineralization and greenhouse  
894 gas emissions in Canadian Cryosols. *Geoderma* 189(189–190): 469–479.
- 895 Parker TC, Subke JA and Wookey PA (2015) Rapid carbon turnover beneath shrub and tree  
896 vegetation is associated with low soil carbon stocks at a subarctic treeline. *Global*  
897 *Change Biology* 21(5): 2070–2081.
- 898 Parmentier F-JW, Christensen TR, Rysgaard S, et al. (2017) A synthesis of the arctic  
899 terrestrial and marine carbon cycles under pressure from a dwindling cryosphere.  
900 *Ambio*, Springer 46: 53–69.
- 901 Pearson RG, Phillips SJ, Lorant MM, et al. (2013) Shifts in Arctic vegetation and associated  
902 feedbacks under climate change. *Nature Climate Change* 3: 673–677.
- 903 Ping CL, Jastrow JD, Jorgenson MT, et al. (2015) Permafrost soils and carbon cycling. 1:  
904 147–171.
- 905 Post E, Forchhammer MC, Bret-Harte MS, et al. (2009) Ecological Dynamics Across the  
906 Arctic Associated with Recent Climate Change. *Science* 325(5946): 1355–1358.
- 907 Poyatos R, Heinemeyer A, Ineson P, et al. (2014) Environmental and Vegetation Drivers of  
908 Seasonal CO<sub>2</sub> Fluxes in a Sub-arctic Forest-Mire Ecotone. *Ecosystems* 17(3): 377–393.



1  
2  
3 909 Raynolds MK, Comiso JC, Walker DA, et al. (2008) Relationship between satellite-derived  
4 910 land surface temperatures, arctic vegetation types, and NDVI. *Remote Sensing of*  
5 911 *Environment* 112(4): 1884–1894.  
6  
7 912 Scherrer D and Körner C (2011) Topographically controlled thermal-habitat differentiation  
8 913 buffers alpine plant diversity against climate warming. *Journal of Biogeography* 38(2):  
9 914 406–416.  
10  
11 915 Schuur EAG, Bockheim J, Canadell JG, et al. (2008) Vulnerability of Permafrost Carbon to  
12 916 Climate Change: Implications for the Global Carbon Cycle. *BioScience*, American  
13 917 Institute of Biological Sciences 58(8): 701.  
14  
15 918 Schuur EAG, Vogel JG, Crummer KG, et al. (2009) The effect of permafrost thaw on old  
16 919 carbon release and net carbon exchange from tundra. *Nature* 459(7246): 556–559.  
17  
18 920 Segal AD and Sullivan PF (2014) Identifying the sources and uncertainties of ecosystem  
19 921 respiration in Arctic tussock tundra. *Biogeochemistry* 121(3): 489–503.  
20  
21 922 Sharp ED, Sullivan PF, Steltzer H, et al. (2013) Complex carbon cycle responses to multi-  
22 923 level warming and supplemental summer rain in the high Arctic. *Global Change*  
23 924 *Biology* 19(6): 1780–1792.  
24  
25 925 Shaver GR, Johnson LC, Cades DH, et al. (1998) Biomass and CO<sub>2</sub> flux in wet sedge tundras  
26 926 responses to nutrients, temperature and light. *Ecological Monographs* 68(1): 75–97.  
27  
28 927 Shaver GR, Street LE, Rastetter EB, et al. (2007) Functional convergence in regulation of net  
29 928 CO<sub>2</sub> flux in heterogeneous tundra landscapes in Alaska and Sweden. *Journal of*  
30 929 *Ecology* 95(4): 802–817.  
31  
32 930 Shaver GR, Rastetter EB, Salmon V, et al. (2013) Pan-Arctic modelling of net ecosystem  
33 931 exchange of CO<sub>2</sub>. *Philosophical Transactions of the Royal Society of London B:*  
34 932 *Biological Sciences* 368(1624): 1–13.  
35  
36 933 Sitch S, McGuire AD, Kimball J, et al. (2007) Assessing the carbon balance of circumpolar  
37 934 arctic tundra using remote sensing and process modeling. *Ecological Applications*  
38 935 17(1): 213–234.  
39  
40 936 Sjögersten S, van der Wal R, Loonen MJJE, et al. (2011) Recovery of ecosystem carbon  
41 937 fluxes and storage from herbivory. *Biogeochemistry* 106(3): 357–370.  
42  
43 938 Smol JP and Douglas MS V (2007) Crossing the final ecological threshold in high Arctic  
44 939 ponds. *PNAS* 104(30): 12395–12397.  
45  
46 940 Stewart KJ, Brummell ME, Coxson DS, et al. (2013) How is nitrogen fixation in the high  
47 941 arctic linked to greenhouse gas emissions? *Plant and Soil* 362(1–2): 215–229.  
48  
49 942 Stocker TF, Qin D, Plattner G-K, et al. (2013) Climate Change 2013 The Physical Science  
50 943 Basis Working Group I Contribution to the Fifth Assessment Report of the  
51 944 Intergovernmental Panel on Climate Change Edited by.  
52  
53 945 Street LE, Shaver GR, Williams M, et al. (2007) What is the relationship between changes in  
54 946 canopy leaf area and changes in photosynthetic CO<sub>2</sub> flux in arctic ecosystems? *Journal*  
55 947 *of Ecology* 95(1): 139–150.  
56  
57 948 Street Lorna E., Shaver GR, Rastetter EB, et al. (2012) Incident radiation and the allocation  
58 949 of nitrogen within Arctic plant canopies: Implications for predicting gross primary  
59 950 productivity. *Global Change Biology* 18(9): 2838–2852.  
60 951 Street L E, Stoy PC, Sommerkorn M, et al. (2012) Seasonal bryophyte productivity in the

- sub-Arctic: a comparison with vascular plants. *Functional Ecology* 26(2): 365–378.
- Susiluoto S, Rasilo T, Pumpanen J, et al. (2008) Effects of grazing on the vegetation structure and carbon dioxide exchange of a fennoscandian fell ecosystem. *Arctic Antarctic and Alpine Research* 40(2): 422–431.
- Tape K, Sturm M and Racine C (2006) The evidence for shrub expansion in Northern Alaska and the Pan-Arctic. *Global Change Biology* 12(4): 686–702.
- Tarnocai C, Canadell JG, Schuur EAG, et al. (2009) Soil organic carbon pools in the northern circumpolar permafrost region. *Global Biogeochemical Cycles* 23(2): 1–11.
- Thomazini A, Francelino MR, Pereira AB, et al. (2016) Geospatial variability of soil CO<sub>2</sub>-C exchange in the main terrestrial ecosystems of Keller Peninsula, Maritime Antarctica. *Science of the Total Environment* 562: 802–811.
- van der Molen MK, van Huissteden J, Parmentier FJW, et al. (2007) The growing season greenhouse gas balance of a continental tundra site in the Indigirka lowlands, NE Siberia. *Biogeosciences* 4: 985–1003.
- van Huissteden J and Dolman AJ (2012) Soil carbon in the Arctic and the permafrost carbon feedback. *Current Opinion in Environmental Sustainability* 4: 545–551.
- Virtanen R, Oksanen L, Oksanen T, et al. (2016) Where do the treeless tundra areas of northern highlands fit in the global biome system: toward an ecologically natural subdivision of the tundra biome. *Ecology and Evolution* 6(1): 143–158.
- Virtanen T and Ek M (2014) The fragmented nature of tundra landscape. *International Journal of Applied Earth Observation and Geoinformation* 27: 4–12.
- Vogel J, Schuur EAG, Trucco C, et al. (2009) Response of CO<sub>2</sub> exchange in a tussock tundra ecosystem to permafrost thaw and thermokarst development. *Journal of Geophysical Research-Biogeosciences* 114(G4): 1–14.
- Vogel Jason, Schuur EAG, Trucco C, et al. (2009) Response of CO<sub>2</sub> exchange in a tussock tundra ecosystem to permafrost thaw and thermokarst development. *Journal of Geophysical Research* 114(G4): G04018.
- Voigt C, Marushchak ME, Lamprecht RE, et al. (2017) Increased nitrous oxide emissions from Arctic peatlands after permafrost thaw. *PNAS* 114(24): 6238–6243.
- Vourlitis George L., Oechel WC, Hope A, et al. (2000) Physiological models for scaling plot measurements of CO<sub>2</sub> flux across an arctic tundra landscape. *Ecological Applications* 10(1): 60–72.
- Vourlitis G L, Oechel WC, Hope A, et al. (2000) Physiological models for scaling plot measurements of CO<sub>2</sub> flux across an arctic tundra landscape. *Ecological Applications* 10(1): 60–72.
- Walker DA, Raynolds MK, Daniels FJA, et al. (2005) The Circumpolar Arctic vegetation map. *Journal of Vegetation Science* 16: 267–282.
- Walker DA, Daniëls FJA, Alsos I, et al. (2016) Circumpolar Arctic vegetation: a hierarchic review and roadmap toward an internationally consistent approach to survey, archive and classify tundra plot data. *Environmental Research Letters* 11(5): 55005.
- Wang Y, Fu B, Lü Y, et al. (2010) Local-scale spatial variability of soil organic carbon and its stock in the hilly area of the Loess Plateau, China. *Quaternary Research* 73(1): 70–76.
- Warren CR and Taranto MT (2011) Ecosystem Respiration in a Seasonally Snow-Covered

1  
2  
3 995 Subalpine Grassland. *Arctic Antarctic and Alpine Research* 43(1): 137–146.  
4 996 Welker JM, Fahnestock JT and Jones MH (2000) Annual CO<sub>2</sub> flux in dry and moist arctic  
5 997 tundra: Field responses to increases in summer temperatures and winter snow depth.  
6 998 *Climatic Change* 44(1–2): 139–150.  
7 999 Welker JM, Fahnestock JT, Henry GHR, et al. (2004) CO<sub>2</sub> exchange in three Canadian High  
8 1000 Arctic ecosystems: Response to long-term experimental warming. *Global Change*  
9 1001 *Biology* 10(12): 1981–1995.  
10 1002 White DM, Garland DS, Ping CL, et al. (2004) Characterizing soil organic matter quality in  
11 1003 arctic soil by cover type and depth. *Cold Regions Science and Technology* 38(1): 63–73.  
12 1004 Williams M, Street LE, van Wijk MT, et al. (2006) Identifying differences in carbon exchange  
13 1005 among arctic ecosystem types. *Ecosystems* 9(2): 288–304.  
14 1006 Xue K, Yuan MM, Shi ZJ, et al. (2016) Tundra soil carbon is vulnerable to rapid microbial  
15 1007 decomposition under climate warming. *Nature Climate Change* 6: 595–600.  
16 1008 Ylänne H, Stark S, Tolvanen A, et al. (2015) Vegetation shift from deciduous to evergreen  
17 1009 dwarf shrubs in response to selective herbivory offsets carbon losses: evidence from  
18 1010 19years of warming and simulated herbivory in the subarctic tundra. *Global Change*  
19 1011 *Biology* 21(10): 3696–3711.  
20 1012 Ylänne H, Stark S and Tolvanen A (2015) Vegetation shift from deciduous to evergreen  
21 1013 dwarf shrubs in response to selective herbivory offsets carbon losses: Evidence from  
22 1014 19 years of warming and simulated herbivory in the subarctic tundra. *Global Change*  
23 1015 *Biology*.  
24 1016 Zamolodchikov DG (2015) CO<sub>2</sub>-exchange in tundra ecosystems of Vaygach Island during  
25 1017 the unusually warm and dry vegetation season. *Zhurnal Obshchei Biologii* 76(2): 83–98.  
26 1018 Zhu RB, Bao T, Wang Q, et al. (2014) Summertime CO<sub>2</sub> fluxes and ecosystem respiration  
27 1019 from marine animal colony tundra in maritime Antarctica. *Atmospheric Environment*  
28 1020 98: 190–201.  
29 1021  
30  
31  
32  
33  
34  
35  
36  
37  
38 1022  
39  
40 1023  
41  
42  
43 1024  
44  
45 1025  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




Figure 1. CO<sub>2</sub> fluxes are regulated by biota, resources, regulators, and disturbance, and vary depending on CAVM (Circumpolar Arctic Vegetation Map) categories resulting in different carbon balances. CAVM represents only the major vegetation units and is developed for landscape or circumpolar scales, although in reality most tundra landscapes are mosaics of several of these types and their subtypes (Walker et al., 2016). The GPP (gross primary production) and ER (ecosystem respiration) bars represent their rates: the darker the shade, the faster the process. Increased GPP is often linked to higher plant respiration rates and, therefore, increased ER (Johnson et al., 2000). Increased GPP also stimulates the rate of litter production, resulting in higher heterotrophic respiration rates (Boelman et al., 2003).

Figure 2. A) The spatial distribution of the observations in the study with a base map of soil organic carbon content. B) The environmental conditions of the observations (in black) versus the environmental conditions in the Arctic tundra (in grey). The data acquired from WorldClimv2 (1970–2000), Soilgrid (topsoil pH), NCSCDv2 (soil organic carbon, SOC), TERRA/MODIS NDVI -product (July, 2015), and GTOPO DEM (elevation) (see S1). The arctic area was defined by Ecoregions (Dinerstein et al., 2017). See S2 for more details.


Figure 3. The hierarchy and relationships of the environmental drivers regulating GPP (gross primary production), ER (ecosystem respiration) and NEE (net ecosystem exchange) from large to fine scale. The arrows represent the main direction of the effect, and the colors of the arrows have no meaning. The resource variables have mostly a direct link to GPP and ER, but regulators and disturbance affect the biota, and thus, GPP and ER.

Figure 4. Number and proportion of studies in which each environmental class was used. Climatological variables and biomass were studied the most, whereas other vegetation properties, soils, and disturbance the least.


Figure 5. A) Number and proportion of studies with observations from different CAVM (Circumpolar Arctic Vegetation Map) categories in the Arctic tundra. B) Number and proportion of studies with observations from the most dominant genera. C) Number and proportion of studies with observations from different CAVM categories per most intensively studied regions. The vegetation data is based on the dominant and secondary species informed by the authors, which was then used to classify the vegetation into CAVM categories. Note the differing proportions on the y-axis.




182x162mm (300 x 300 DPI)




178x218mm (300 x 300 DPI)



178x98mm (300 x 300 DPI)



81x80mm (300 x 300 DPI)



178x130mm (300 x 300 DPI)

Table 1. Environmental categories and classes used to explain CO<sub>2</sub> fluxes. Leaf area index (LAI) measures the amount of leaf area in a canopy per unit ground area (m<sup>2</sup> m<sup>-2</sup>; Asner et al., 2003). The normalized difference vegetation index (NDVI) or some other indices derived from remote sensing describe the amount of green biomass and/or LAI. Number in parentheses indicate the number of studies (0-93) to include the environmental categories/classes.

Categories	Climate (91)	Vegetation (79)	Soils (54)	Disturbance (38)
Classes	Air temperature (80)  Radiation (72) - Photosynthetically active radiation (PAR)	Biomass (75) -cover, biomass, LAI -remotely sensed parameters (NDVI)  Plant traits (35) -leaf N -leaf cover/area	Soil nutrients (36) -C, N, P, -pH -litter biomass  Physical properties (41) -soil type, texture	Geomorphology (24) -permafrost thaw, cryoturbation, fluvial processes  Herbivory (13) -small & large mammals



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

	Soil microclimate (81) -topsoil temperature -active layer depth -moisture -water table height	-vegetation height  Growth & phenology (17) -shoot, root, leaf, biomass growth -phenology	-organic layer depth  Microbes (8) -microbial biomass, activity, community structure	-birds -insect outbreaks  Disease (1) -plant pathogens
--	--	---	---	--

For Peer Review

Category	Knowledge level	Future research needs	Mechanisms	Importance
<b>Fluxes</b>	Good	Combination of soil respiration and NEE.  Linkages to other greenhouse gas fluxes.  Upscaling of fluxes.	Soil respiration & ER define NEE in a changing climate and disturbance regime.  The role of other gas fluxes in global change.  More detailed information on spatio-temporal variation and hot spots.	Urgent
<b>Climate</b>	Very good	Temperature extremes.  Growing-season length.	Climate warming and changes in precipitation.  Changes in dynamics.	Less urgent
<b>Vegetation</b>	Good	Barren and low-shrubs.  Cryptogams.	Shrubification, vegetation shift.  The role of mosses.	Moderate
<b>Soils</b>	Poor	Soil nutrients, SOC stocks and fluxes.  Microbial biomass, community and activity.	Substrate quality important for GPP, regulates temperature sensitivity of ER. Changes in SOC stocks.  Microbial effects and response to global warming.	Urgent
<b>Disturbance</b>	Poor	Geomorphology, permafrost dynamics, fire, disease.	Potential changes in geomorphological processes. Increase of fire and disease outbreaks. More CO <sub>2</sub> and fire to the	Urgent

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

			atmosphere.	
		Herbivory.	Herbivory effects of different species in various environments.	

Table 2. Chamber-derived CO<sub>2</sub> fluxes and environmental categories studied, their knowledge level, research gaps, and importance of the topics in the global change - context.

For Peer Review

## Appendix

### Supplementary Material 1. Articles in the review.

- Arens SJT, Sullivan PF and Welker JM (2008) Nonlinear responses to nitrogen and strong interactions with nitrogen and phosphorus additions drastically alter the structure and function of a high arctic ecosystem. *Journal of Geophysical Research-Biogeosciences* 113(G3): 10.
- Arndal MF, Illeris L, Michelsen A, et al. (2009) Seasonal Variation in Gross Ecosystem Production, Plant Biomass, and Carbon and Nitrogen Pools in Five High Arctic Vegetation Types. *Arctic Antarctic and Alpine Research* 41(2): 164–173.
- Biasi C, Meyer H, Rusalimova O, et al. (2008) Initial effects of experimental warming on carbon exchange rates, plant growth and microbial dynamics of a lichen-rich dwarf shrub tundra in Siberia. *Plant and Soil* 307(1–2): 191–205.
- Boelman NT, Stieglitz M, Rueth HM, et al. (2003) Response of NDVI, biomass, and ecosystem gas exchange to long-term warming and fertilization in wet sedge tundra. *Oecologia* 135(3): 414–421.
- Bokhorst S, Bjerke JW, Street LE, et al. (2011) Impacts of multiple extreme winter warming events on sub-Arctic heathland: phenology, reproduction, growth, and CO<sub>2</sub> flux responses. *Global Change Biology* 17(9): 2817–2830.
- Bokhorst S, Phoenix GK, Berg MP, et al. (2015) Climatic and biotic extreme events moderate long-term responses of above- and belowground sub-Arctic heathland communities to climate change. *Global Change Biology* 21(11): 4063–4075.
- Brummell ME, Farrell RE and Siciliano SD (2012) Greenhouse gas soil production and surface fluxes at a high arctic polar oasis. *Soil Biology and Biochemistry* 52: 1–12.
- Bäckstrand K, Crill PM, Jackowicz-Korcz Nski M, et al. (2010) Annual carbon gas budget for a subarctic peatland, Northern Sweden. *Biogeosciences* 7(1): 95–108.
- Cahoon SMP, Sullivan PF and Post E (2016) Greater Abundance of *Betula nana* and Early Onset of the Growing Season Increase Ecosystem CO<sub>2</sub> Uptake in West Greenland. *Ecosystems* 19(7): 1149–1163.
- Cahoon SMP, Sullivan PF, Post E, et al. (2012) Large herbivores limit CO<sub>2</sub> uptake and suppress carbon cycle responses to warming in West Greenland. *Global Change Biology* 18(2): 469–479.

Cahoon SMP, Sullivan PF, Shaver GR, et al. (2012) Interactions among shrub cover and the soil microclimate may determine future Arctic carbon budgets. *Ecology Letters* 15(12): 1415–1422.

Christensen TR, Friborg T, Sommerkorn M, et al. (2000) Trace gas exchange in a high-arctic valley 1. Variations in CO<sub>2</sub> and CH<sub>4</sub> flux between tundra vegetation types. *Global Biogeochemical Cycles* 14(3): 701–713.

Curasi SR, Loranty MM and Natali SM (2016) Water track distribution and effects on carbon dioxide flux in an eastern Siberian upland tundra landscape. *Environmental Research Letters* 11(4): 45002.

Dagg J and Lafleur P (2010) An application of plot-scale NDVI in predicting carbon dioxide exchange and leaf area index in heterogeneous subarctic tundra. *Canadian Journal of Remote Sensing* 36: S111–S123.

Dagg J and Lafleur P (2011) Vegetation Community, Foliar Nitrogen, and Temperature Effects on Tundra CO<sub>2</sub> Exchange across a Soil Moisture Gradient. *Arctic Antarctic and Alpine Research* 43(2): 189–197.

Davidson SJ, Sloan VL, Phoenix GK, et al. (2016) Vegetation Type Dominates the Spatial Variability in CH<sub>4</sub> Emissions Across Multiple Arctic Tundra Landscapes. *Ecosystems* 19(6): 1116–1132.

Douma JC, Van Wijk MT, Lang SI, et al. (2007) The contribution of mosses to the carbon and water exchange of arctic ecosystems: quantification and relationships with system properties. *Plant Cell and Environment* 30(10): 1205–1215.

Falk JM, Schmidt NM and Ström L (2014) Effects of simulated increased grazing on carbon allocation patterns in a high arctic mire. *Biogeochemistry* 119(1–3): 229–244.

Falk JM, Schmidt NM, Christensen TR, et al. (2015) Large herbivore grazing affects the vegetation structure and greenhouse gas balance in a high arctic mire. *Environmental Research Letters* 10(4): 11.

Fox AM, Huntley B, Lloyd CR, et al. (2008) Net ecosystem exchange over heterogeneous Arctic tundra: Scaling between chamber and eddy covariance measurements. *Global Biogeochemical Cycles* 22(2): 1–15.

Griffis TJ, Rouse WR and Waddington JM (2000) Scaling net ecosystem CO<sub>2</sub> exchange from the community to landscape-level at a subarctic fen. *Global Change Biology* 6(4): 459–473.



- Grogan P and Chapin FS (2000) Initial effects of experimental warming on above- and belowground components of net ecosystem CO<sub>2</sub> exchange in arctic tundra. *Oecologia* 125(4): 512–520.
- Grogan P and Jonasson S (2005) Temperature and substrate controls on intra-annual variation in ecosystem respiration in two subarctic vegetation types. *Global Change Biology* 11(3): 465–475.
- Heikkinen JEP, Elsakov V and Martikainen PJ (2002) Carbon dioxide and methane dynamics and annual carbon balance in tundra wetland in NE Europe, Russia. *Global Biogeochemical Cycles* 16(4): 16.
- Heikkinen JEP, Maljanen M, Aurela M, et al. (2002) Carbon dioxide and methane dynamics in a sub-Arctic peatland in northern Finland. *Polar Research* 21(1): 49–62.
- Heikkinen JEP, Virtanen T, Huttunen JT, et al. (2004) Carbon balance in East European tundra. *Global Biogeochemical Cycles* 18(1): 1–14.
- Huemmrich KF, Kinoshita G, Gamon JA, et al. (2010) Tundra carbon balance under varying temperature and moisture regimes. *Journal of Geophysical Research-Biogeosciences* 115(G4): 1–8.
- Illeris L, König SM, Grogan P, et al. (2004) Growing-season carbon dioxide flux in a dry subarctic heath: Responses to long-term manipulations. *Arctic Antarctic and Alpine Research* 36(4): 456–463.
- Johansson T, Malmer N, Crill PM, et al. (2006) Decadal vegetation changes in a northern peatland, greenhouse gas fluxes and net radiative forcing. *Global Change Biology* 12(12): 2352–2369.
- Johnson LC, Shaver GR, Cades DH, et al. (2000) Plant carbon-nutrient interactions control CO<sub>2</sub> exchange in Alaskan wet sedge tundra ecosystems. *Ecology* 81(2): 453–469.
- Kade A, Bret-Harte MS, Euskirchen ES, et al. (2012) Upscaling of CO<sub>2</sub> fluxes from heterogeneous tundra plant communities in Arctic Alaska. *Journal of Geophysical Research - Biogeosciences* 117(G4): G04007.
- Karelin D V., Zamolodchikov DG, Zukert N V., et al. (2013) Interannual changes in PAR and soil moisture during the warm season may be more important for directing of annual carbon balance in tundra than temperature fluctuations. *Zhurnal Obshchei Biologii* 74(1): 3–22.
- Kelsey KC, Leffler AJ, Beard KH, et al. (2016) Interactions among climate, topography and herbivory control greenhouse gas (CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O) fluxes in a subarctic coastal wetland. *Journal of Geophysical Research - Biogeosciences* 121(12): 2960–2975.

Kim Y, Lee B-Y, Suzuki R, et al. (2016) Spatial characteristics of ecosystem respiration in three tundra ecosystems of Alaska. *Polar Science* 10(3): 356–363.

Kim Y, Nishina K, Chae N, et al. (2014) Constraint of soil moisture on CO<sub>2</sub> efflux from tundra lichen, moss, and tussock in Council, Alaska, using a hierarchical Bayesian model. *Biogeosciences* 11: 5567–5579.

Kutzbach L, Schneider J, Sachs T, et al. (2007) CO<sub>2</sub> flux determination by closed-chamber methods can be seriously biased by inappropriate application of linear regression. *Biogeosciences* 4: 1005–1025.

Kwon MJ, Heimann M, Kolle O, et al. (2016) Long-term drainage reduces CO<sub>2</sub> uptake and increases CO<sub>2</sub> emission on a Siberian floodplain due to shifts in vegetation community and soil thermal characteristics. *Biogeosciences* 13: 4219–4235.

La Puma IP, Philippi TE and Oberbauer SF (2007) Relating NDVI to ecosystem CO<sub>2</sub> exchange patterns in response to season length and soil warming manipulations in arctic Alaska. *Remote Sensing of Environment* 109(2): 225–236.

Lara MJ, McGuire AD, Euskirchen ES, et al. (2015) Polygonal tundra geomorphological change in response to warming alters future CO<sub>2</sub> and CH<sub>4</sub> flux on the Barrow Peninsula. *Global Change Biology* 21(4): 1634–1651.

Lara MJ, Villarreal S, Johnson DR, et al. (2012) Estimated change in tundra ecosystem function near Barrow, Alaska between 1972 and 2010. *Environmental Research Letters* 7(1): 15507.

Larsen KS, Ibrom A, Jonasson S, et al. (2007) Significance of cold-season respiration and photosynthesis in a subarctic heath ecosystem in Northern Sweden. *Global Change Biology* 13(7): 1498–1508.

Leffler AJ, Klein ES, Oberbauer SF, et al. (2016) Coupled long-term summer warming and deeper snow alters species composition and stimulates gross primary productivity in tussock tundra. *Oecologia* 181(1): 287–297.

López-Blanco E, Lund M, Williams M, et al. (2016) Exchange of CO<sub>2</sub> in Arctic tundra: impacts of meteorological variations and biological disturbance. *Biogeosciences Discussions* doi:10.519.

Lund M, Christensen TR, Mastepanov M, et al. (2009) Effects of N and P fertilization on the greenhouse gas exchange in two northern peatlands with contrasting N deposition rates. *Biogeosciences* 6(10): 2135–2144.

Maanavilja L, Riutta T, Aurela M, et al. (2011) Spatial variation in CO<sub>2</sub> exchange at a northern aapa mire. *Biogeochemistry* 104(1–3): 325–345.

- 1  
2  
3 Malhotra A and Roulet NT (2015) Environmental correlates of peatland carbon fluxes in a  
4 thawing landscape: do transitional thaw stages matter? *Biogeosciences* 12(10): 3119–  
5 3130.  
6  
7  
8 Marushchak ME, Kiepe I, Biasi C, et al. (2013) Carbon dioxide balance of subarctic tundra  
9 from plot to regional scales. *Biogeosciences* 10(1): 437–452.  
10  
11 Mastepanov M, Sigsgaard C, Tagesson T, et al. (2013) Revisiting factors controlling  
12 methane emissions from high-Arctic tundra. *Biogeosciences* 10(11): 5139–5158.  
13  
14  
15 McEwing KR, Fisher JP and Zona D (2015) Environmental and vegetation controls on the  
16 spatial variability of CH<sub>4</sub> emission from wet-sedge and tussock tundra ecosystems in  
17 the Arctic. *Plant and Soil* 388(1–2): 37–52.  
18  
19  
20 Metcalfe DB and Olofsson J (2015) Distinct impacts of different mammalian herbivore  
21 assemblages on arctic tundra CO<sub>2</sub> exchange during the peak of the growing season.  
22 *Oikos* 124(12): 1632–1638.  
23  
24  
25 Natali SM, Schuur EAG, Trucco C, et al. (2011) Effects of experimental warming of air, soil  
26 and permafrost on carbon balance in Alaskan tundra. *Global Change Biology* 17(3):  
27 1394–1407.  
28  
29  
30 Nobrega S and Grogan P (2008) Landscape and ecosystem-level controls on net carbon  
31 dioxide exchange along a natural moisture gradient in Canadian low arctic tundra.  
32 *Ecosystems* 11(3): 377–396.  
33  
34  
35 Nykanen H, Heikkinen JEP, Pirinen L, et al. (2003) Annual CO<sub>2</sub> exchange and CH<sub>4</sub> fluxes on  
36 a subarctic palsamire during climatically different years. *Global Biogeochemical Cycles*  
37 17(1): 18.  
38  
39  
40 Oberbauer SF, Tweedie CE, Welker JM, et al. (2007) Tundra CO<sub>2</sub> fluxes in response to  
41 experimental warming across latitudinal and moisture gradients. *Ecological*  
42 *Monographs* 77(2): 221–238.  
43  
44  
45 Olivas PC, Oberbauer SF, Tweedie C, et al. (2011) Flux Components of Alaskan Coastal Plain  
46 Tundra: Response to Contrasting Growing Seasons. *Arctic, Antarctic, and Alpine*  
47 *Research* 43(2): 256–266.  
48  
49  
50 Olivas PC, Oberbauer SF, Tweedie CE, et al. (2010) Responses of CO<sub>2</sub> flux components of  
51 Alaskan Coastal Plain tundra to shifts in water table. *Journal of Geophysical Research-*  
52 *Biogeosciences* 115(G4): 1–13.  
53  
54  
55 Olofsson J, Ericson L, Torp M, et al. (2011) Carbon balance of Arctic tundra under increased  
56 snow cover mediated by a plant pathogen. *Nature Climate Change* 1(4): 220–223.  
57  
58  
59  
60

Poyatos R, Heinemeyer A, Ineson P, et al. (2014) Environmental and Vegetation Drivers of Seasonal CO<sub>2</sub> Fluxes in a Sub-arctic Forest-Mire Ecotone. *Ecosystems* 17(3): 377–393.

Rogers MC, Sullivan PF and Welker JM (2011) Evidence of nonlinearity in the response of net ecosystem CO<sub>2</sub> exchange to increasing levels of winter snow depth in the high Arctic of northwest Greenland. *Arctic, Antarctic, and Alpine Research* 43(1): 95–106.

Schuur EAG, Vogel JG, Crummer KG, et al. (2009) The effect of permafrost thaw on old carbon release and net carbon exchange from tundra. *Nature* 459(7246): 556–559.

Segal AD and Sullivan PF (2014) Identifying the sources and uncertainties of ecosystem respiration in Arctic tussock tundra. *Biogeochemistry* 121(3): 489–503.

Sharp ED, Sullivan PF, Steltzer H, et al. (2013) Complex carbon cycle responses to multi-level warming and supplemental summer rain in the high Arctic. *Global Change Biology* 19(6): 1780–1792.

Shaver GR, Rastetter EB, Salmon V, et al. (2013) Pan-Arctic modelling of net ecosystem exchange of CO<sub>2</sub>. *Philosophical Transactions of the Royal Society of London B: Biological Sciences* 368(1624): 1–13.

Shaver GR, Street LE, Rastetter EB, et al. (2007) Functional convergence in regulation of net CO<sub>2</sub> flux in heterogeneous tundra landscapes in Alaska and Sweden. *Journal of Ecology* 95(4): 802–817.

Sjögersten S, Van Der Wal R and Woodin SJ (2008) Habitat type determines herbivory controls over CO<sub>2</sub> fluxes in a warmer arctic. *Ecology* 89(8): 2103–2116.

Sjögersten S, Wal R van der, Loonen MJJE, et al. (n.d.) Recovery of ecosystem carbon fluxes and storage from herbivory. *Biogeochemistry*, Springer Netherlands 106(3): 357–370.

Sjögersten S, Van Der R, Ae W, et al. (2006) Small-scale hydrological variation determines landscape CO<sub>2</sub> fluxes in the high Arctic. *Biogeochemistry* 80: 205–216.

Soegaard H, Nordstroem C, Friberg T, et al. (2000) Trace gas exchange in a high-Arctic valley: 3. Integrating and scaling CO<sub>2</sub> fluxes from canopy to landscape using flux data, footprint modeling, and remote sensing. *Global Biogeochemical Cycles* 14(3): 725–744.

Stewart KJ, Brummell ME, Coxson DS, et al. (2013) How is nitrogen fixation in the high arctic linked to greenhouse gas emissions? *Plant and Soil* 362(1–2): 215–229.

Street LE, Shaver GR, Rastetter EB, et al. (2012) Incident radiation and the allocation of nitrogen within Arctic plant canopies: Implications for predicting gross primary productivity. *Global Change Biology* 18(9): 2838–2852.

- Street LE, Shaver GR, Williams M, et al. (2007) What is the relationship between changes in canopy leaf area and changes in photosynthetic CO<sub>2</sub> flux in arctic ecosystems? *Journal of Ecology* 95(1): 139–150.
- Street LE, Stoy PC, Sommerkorn M, et al. (2012) Seasonal bryophyte productivity in the sub-Arctic: a comparison with vascular plants. *Functional Ecology* 26(2): 365–378.
- Strom L, Tagesson T, Mastepanov M, et al. (2012) Presence of *Eriophorum scheuchzeri* enhances substrate availability and methane emission in an Arctic wetland. *Soil Biology & Biochemistry* 45: 61–70.
- Sullivan PF, Arens SJT, Chimner RA, et al. (2008) Temperature and microtopography interact to control carbon cycling in a high arctic fen. *Ecosystems* 11(1): 61–76.
- Susiluoto S, Rasilo T, Pumpanen J, et al. (2008) Effects of grazing on the vegetation structure and carbon dioxide exchange of a fennoscandian fell ecosystem. *Arctic Antarctic and Alpine Research* 40(2): 422–431.
- Tagesson T, Mastepanov M, Tamstorf MP, et al. (2010) Increased GPP in a high arctic fen 1992–2008 Satellites reveal an increase in gross primary production in a greenlandic high arctic fen 1992–2008 Increased GPP in a high arctic fen 1992–2008. *Biogeosciences Discussion* 7: 1101–1129.
- Trucco C, Schuur EAG, Natali SM, et al. (2012) Seven-year trends of CO<sub>2</sub> exchange in a tundra ecosystem affected by long-term permafrost thaw. *Journal of Geophysical Research: Biogeosciences* 117(G2): 1–12.
- Uchida M, Kishimoto A, Muraoka H, et al. (2010) Seasonal shift in factors controlling net ecosystem production in a high Arctic terrestrial ecosystem. *Journal of Plant Research* 123(1): 79–85.
- van der Molen MK, van Huissteden J, Parmentier FJW, et al. (2007) The growing season greenhouse gas balance of a continental tundra site in the Indigirka lowlands, NE Siberia. *Biogeosciences* 4: 985–1003.
- van Der Wal R, Sjögersten S, Woodin SJ, et al. (2007) Spring feeding by pink-footed geese reduces carbon stocks and sink strength in tundra ecosystems. *Global Change Biology* 13(2): 539–545.
- Welker JM, Fahnestock JT and Jones MH (2000) Annual CO<sub>2</sub> flux in dry and moist arctic tundra: Field responses to increases in summer temperatures and winter snow depth. *Climatic Change* 44(1–2): 139–150.



Welker JM, Fahnestock JT, Henry GHR, et al. (2004) CO<sub>2</sub> exchange in three Canadian High Arctic ecosystems: response to long-term experimental warming. *Global Change Biology* 10(12): 1981–1995.

Williams M, Rastetter EB, Van der Pol L, et al. (2014) Arctic canopy photosynthetic efficiency enhanced under diffuse light, linked to a reduction in the fraction of the canopy in deep shade. *New Phytologist* 202(4): 1266–1276.

Williams M, Street LE, van Wijk MT, et al. (2006) Identifying differences in carbon exchange among arctic ecosystem types. *Ecosystems* 9(2): 288–304.

Vogel J, Schuur EAG, Trucco C, et al. (2009) Response of CO<sub>2</sub> exchange in a tussock tundra ecosystem to permafrost thaw and thermokarst development. *Journal of Geophysical Research-Biogeosciences* 114(G4): 1–14.

Vourlitis GL, Oechel WC, Hope A, et al. (2000) Physiological models for scaling plot measurements of CO<sub>2</sub> flux across an arctic tundra landscape. *Ecological Applications* 10(1): 60–72.

Väisänen M, Yläne H, Kaarlejärvi E, et al. (2014) Consequences of warming on tundra carbon balance determined by reindeer grazing history. *Nature Climate Change* 4(5): 384–388.

Yläne H, Stark S, Tolvanen A, et al. (2015) Vegetation shift from deciduous to evergreen dwarf shrubs in response to selective herbivory offsets carbon losses: evidence from 19 years of warming and simulated herbivory in the subarctic tundra. *Global Change Biology* 21(10): 3696–3711.

Zamolodchikov D, Karelin D and Ivaschenko A (2000) Sensitivity of tundra carbon balance to ambient temperature. *Water, Air, and Soil Pollution* 119: 157–169.

Zamolodchikov DG (2015) CO<sub>2</sub>-exchange in tundra ecosystems of Vaygach Island during the unusually warm and dry vegetation season. *Zhurnal Obshchei Biologii* 76(2): 83–98.

Zamolodchikov DG and Karelin D V (2001) An empirical model of carbon fluxes in Russian tundra. *Global Change Biology* 7(2): 147–161.

Zamolodchikov DG, Karelin D V., Ivaschenko AI, et al. (2003) CO<sub>2</sub> flux measurements in Russian Far East tundra using eddy covariance and closed chamber techniques. *Tellus B* 55(4): 879–892.

Zona D, Lipson DA, Zulueta RC, et al. (2011) Microtopographic controls on ecosystem functioning in the Arctic Coastal Plain. *Journal of Geophysical Research: Biogeosciences* 116(3): 1–12.

## Supplementary Material 2. Methods in the systematic review

### 1. Data collection

This article is based on a literature review of chamber-derived CO<sub>2</sub> flux studies during the growing season in the Arctic tundra. The survey was conducted via ISI Web of Science (WoS) for articles published in the years 2000-2016. The search was carried out using a query that accounted for the region, scale, flux terminology, and different vegetation types: (tundra or arctic) and ecosystem and ("CO<sub>2</sub> flux" or "carbon dioxide emissions" or "greenhouse gas exchange" or "CO<sub>2</sub> exchange" or "carbon exchange" or "carbon flux") and (meadow or sedge or tussock or hummock or heath or herb or grass or grassland or graminoid or forb or moss or bryophyte or lichen or "cushion plant" or shrub or tree). The query resulted in 242 articles out of which we included approximately 20 % of the studies. Firstly, we wanted to focus on soil chamber measurements and, therefore, excluded studies with eddy covariance or leaf cuvette measurements only. Secondly, only studies that included growing season measurements were taken into account. Studies with GPP, ER, and/or NEE measurements were included. Boreal regions were excluded from the review as we wanted to focus on tundra patterns and processes only, although they are usually included in the carbon balance assessments of Arctic regions (Belshe et al., 2013; McGuire et al., 2010; Treat et al., 2015). However, we included studies with sub-Arctic and Arctic peatlands and fens due to their importance in tundra carbon cycling. Additional articles were derived from the references of the selected publications.

The final 93 publications were read through systematically in order to collect data on the 1) measurement information, 2) study design and location, and 3) environmental variables used to explain the CO<sub>2</sub> fluxes. All the environmental variables measured and described were taken into account, also derived from the study design descriptions (e.g. organic layer depth). The variables were further divided into the following categories and their classes: climate (air temperature, photosynthetically active radiation, soil temperature, soil moisture), vegetation (biomass, plant traits, growth & phenology), soils (physical, chemical and biological) and disturbance (geomorphology, herbivory, disease). To our knowledge, at least seven articles were partly using the same CO<sub>2</sub> flux data, but the other environmental variables used in these studies was variable.

Information on the dominant and secondary species was also collected, and the vegetation data was further classified into Circumpolar Arctic Vegetation Map (CAVM) physiognomic units (Walker et al., 2005). CAVM was developed for landscape-scale analysis and does not consider fine-scale variation but we wanted to classify the vegetation data according to a widely-used vegetation classification. Furthermore, we recognize that classification was sometimes subjective and based on limited data on the species cover. Principally, we defined plots that were described to have only limited vegetation cover, or were described

as barren or polar desert, to physiognomic unit B; plots that had tussocks and were mainly dominated by *Eriophorum*, *Carex*, and mosses to unit G; plots with a dominance of *Dryas*, *Cassiope*, and other prostrate-dwarf shrubs, such as *Salix arctica* and *S. polaris*, to unit P; erect-shrubs such as *Betula nana*, *Salix glauca*, *Rhododendron tomentosum*, *Vaccinium vitis-idaea* to unit S; and, plots described to be on wetland (bog, fen) to unit W.

2. Other data sets used in the figures

The data for Figure 4 was acquired from WorldClim 1970-2000 (average temperature and yearly cumulative precipitation at 2.5 minutes resolution; Fick & Hijmans, 2017), Soilgrid (topsoil pH at approx. 1 km resolution; Hengl et al., 2017), NCSCDv2 (soil organic carbon content at approx. 10 km resolution; Hugelius et al., 2013), TERRA/MODIS NDVI-product (from 1<sup>st</sup> of July, 2015 at approx. 10 km resolution; NASA, 2017) and GTOPO DEM (elevation at approx. 1 km resolution; USGS, 2004). The data was processed with the *raster* package (Hijmans et al. 2016) in R program. The values from the circumpolar data sets were extracted to the observation points. Raster extent varied and, therefore, 10-25 observation sites did not receive any values given they were outside the extent (e.g. soil pH map in southern Svalbard). The tundra area for describing the entire Arctic variation was defined by Ecoregions (Dinerstein et al., 2017). The entire Arctic variation data was rescaled to 10 km resolution for the plots.

Supplementary Material 3. Table with 20 most cited publications. Citations were derived from Web of Science, 16<sup>th</sup> of March 2017. Fluxes describe the CO<sub>2</sub> fluxes measured in the study (SR= soil respiration), manipulation defines whether the study is based on manipulative observations (e.g. experimental warming) and region defines study location(s) of the observations.

Authors	Year	Title	Journal	Citations	Fluxes	Manipulation	Region
Schuur et al.	2009	The effect of permafrost thaw on old carbon release and net carbon exchange from tundra	NATURE	423	GPP, ER, NEE	0	Alaska
Welker et al.	2000	Annual CO <sub>2</sub> flux in dry and moist arctic tundra: Field responses to increases in summer temperatures and winter snow depth	CLIMATIC CHANGE	131	NEE	1	Alaska
Oberbauer et al.	2007	Tundra CO <sub>2</sub> fluxes in response to experimental warming across latitudinal and moisture gradients	ECOLOGICAL MONOGRAPHS	124	GPP, ER, NEE	1	Canada
Welker et al.	2004	CO <sub>2</sub> exchange in three Canadian High Arctic ecosystems: response to long-term experimental warming	GLOBAL CHANGE BIOLOGY	117	GPP, ER, NEE	1	Canada
Kutzbach et al.	2007	CO <sub>2</sub> flux determination by closed-chamber methods can be seriously biased by inappropriate application of linear regression	BIOGEOSCIENCES	109	NEE	0	Fennoscandia, Russia
Christensen et al.	2000	Trace gas exchange in a high-arctic valley 1. Variations in CO <sub>2</sub> and CH <sub>4</sub> flux between tundra vegetation types	GLOBAL BIOGEOCHEMICAL CYCL	107	GPP, ER, NEE	0	Greenland
Johansson et al.	2006	Decadal vegetation changes in a northern peatland, greenhouse gas fluxes and net radiative forcing	GLOBAL CHANGE BIOLOGY	106	NEE	1	Fennoscandia
Boelman et al.	2003	Response of NDVI, biomass, and ecosystem gas exchange to long-term warming and fertilization in wet sedge tundra	OECOLOGIA	104	GPP, ER, NEE	1	Alaska
Shaver et al.	2007	Functional convergence in regulation of net CO <sub>2</sub> flux in heterogeneous tundra landscapes in Alaska and Sweden	JOURNAL OF ECOLOGY	92	GPP, ER, NEE	0	Fennoscandia
Steed et al.	2007	What is the relationship between changes in canopy leaf area and changes in photosynthetic CO <sub>2</sub> flux in arctic ecosystems?	JOURNAL OF ECOLOGY	87	GPP	0	Fennoscandia, Alaska
Johnson et al.	2000	Plant carbon-nutrient interactions control CO <sub>2</sub> exchange in Alaskan wet sedge tundra ecosystems	ECOLOGY	81	GPP, ER, NEE, SR	1	Alaska
Hakkinen et al.	2004	Carbon balance in East European tundra	GLOBAL BIOGEOCHEMICAL CYCL	59	GPP, ER, NEE, SR	0	Russia
Biasi et al.	2008	Initial effects of experimental warming on carbon exchange rates, plant growth and microbial dynamics of a lichen-rich dwarf shrub tundra in Siberia	PLANT AND SOIL	59	GPP, ER, NEE	1	Siberia
Bokhorst et al.	2011	Impacts of multiple extreme winter warming events on sub-Arctic heathland: phenology, reproduction, growth, and CO <sub>2</sub> flux responses	GLOBAL CHANGE BIOLOGY	56	GPP, ER, NEE, SR	1	Fennoscandia
Sullivan et al.	2008	Temperature and microtopography interact to control carbon cycling in a high arctic fen	ECOSYSTEMS	54	GPP, ER, NEE	1	Greenland
Grogan & Johnsson	2005	Temperature and substrate controls on intra-annual variation in ecosystem respiration in two subarctic vegetation types	GLOBAL CHANGE BIOLOGY	54	ER	1	Fennoscandia
Grogan & Chapin	2000	Initial effects of experimental warming on above- and belowground components of net ecosystem CO <sub>2</sub> exchange in arctic tundra	OECOLOGIA	54	GPP, ER, NEE, SR	1	Alaska
Natali et al.	2011	Effects of experimental warming of air, soil and permafrost on carbon balance in Alaskan tundra	GLOBAL CHANGE BIOLOGY	53	GPP, ER, NEE, SR	1	Alaska
Backstrand et al.	2010	Annual carbon gas budget for a subarctic peatland, Northern Sweden	BIOGEOSCIENCES	51	NEE	0	Fennoscandia
Larsen et al.	2007	Significance of cold-season respiration and photosynthesis in a subarctic heath ecosystem in Northern Sweden	GLOBAL CHANGE BIOLOGY	47	GPP, ER, NEE	1	Fennoscandia

Supplementary Material 4. Descriptive data from the 93 articles included in this review. The different columns indicate studies with GPP, ER, and/or NEE measurements, and an average of all CO<sub>2</sub> flux studies. The row names are different classifications of the data which describe information of the studies, study designs, methods, measurements, vegetation, and climatological, soil, and disturbance variables. Cells indicate the number of the studies (except for average number of citations). In the spatial study designs, the focus is on understanding the CO<sub>2</sub> fluxes in the landscape (related to e.g. different vegetation types or topographic positions) whereas in the temporal study designs, the focus is to understand CO<sub>2</sub> dynamics. In manipulative studies, environmental conditions have been modified. In some cases, no data was available (e.g. chamber size).

Studies with measurements of		GPP	ER	NEE	All fluxes
General information		Number of	studies with	observations	of
Arctic	Arctic	28	29	29	32
	Oroarctic	31	33	36	39
	High Arctic	27	30	29	33
Region	Alaska	26	29	27	32
	Canada	7	7	7	8
	Fennoscandia	20	22	24	27
	Russia	12	12	13	13
	Greenland	16	15	17	18
	Svalbard	5	7	7	8
Publication year	2010-2016	32	36	34	39
	2005-2010	19	19	20	23
	2000-2005	9	9	9	9
Average number of citations		35	34	37	36
Study designs					
Spatial/temporal	Spatial	21	22	24	25
	Temporal	18	20	19	21
	Spatial and temporal	31	34	33	39

Observational/manipulative	Observational	42	46	47	53
	Manipulative	35	37	36	39
	Warming	20	20	21	21
	Nutrient addition	7	6	6	7
	Water modifications	5	5	5	5
Median number of observations	All	24	24	29	26
	Spatial	40	40	41	42
	Temporal	18	18	18	18
	Spatial and temporal	18	19	18	20
	Manipulative	20	20	20	20
	Observational	30	29	26	29
Median number of sites	All	3	3	3	3
	Observational	4	4	4	4
<b>Study methods</b>					
Chamber types	Manual	70	75	73	82
	Automatic	8	9	11	11
Chamber size	Edges/diameter < 25 cm	39	43	42	46
	Edges/diameter 25-50 cm	22	24	23	26
	Edges/diameter > 50 cm	17	17	19	21
	Height < 25 cm	45	49	46	52
	Height 25-50 cm	17	20	21	23
	Height > 50 cm	9	9	9	9
Chamber type	LI-COR	48	53	51	59
	EGM	15	15	15	16
	Vaisala	2	2	2	2



<b>CO<sub>2</sub> measurements</b>					
Fluxes	GPP	78	76	74	78
	ER	76	84	77	84
	NEE	74	77	84	84
	Soil respiration	12	14	14	15
Temporal measurements	Summer season (J-A)	58	63	62	69
	Cold-season (A, M, S, O)	14	18	20	21
	Winter	16	17	18	19
	Yearly	29	33	33	35
Other flux measurements	Eddy covariance	9	10	11	11
	Leaf cuvette	4	5	4	5
Other gas fluxes	N <sub>2</sub> O	2	2	2	3
	CH <sub>4</sub>	20	21	23	24
<b>Vegetation</b>					
Peatland		21	22	23	23
Vegetation cover	Species/PFT/Total cover	44	44	44	44
Biomass	Aboveground species/PFT/Total	44	43	43	44
	Belowground biomass	10	12	12	12
	LAI	21	21	20	23
	NDVI	27	27	26	29
	(field or satellite-based)				
Any biomass (cover + biomass)		75	75	75	75
Plant traits	Leaf chemical properties	15	15	15	17

	Leaf cover	11	12	12	13
	Vegetation height	9	10	10	11
Growth & phenology		17	17	17	17
<b>Climate</b>					
Surface	Air temperature	74	77	78	80
	PAR	69	70	71	72
Soils	Soil temperature	64	69	66	74
	Thaw depth	37	41	42	45
	Soil moisture	39	42	39	46
	Water table height	32	33	34	35
<b>Soils</b>					
Nutrients	C	15	15	15	15
	N	18	18	18	18
	P	5	5	5	5
	pH	13	13	13	15
	Litter biomass	6	6	5	7
Physical	Organic layer depth	25	28	25	28
	Soil type	13	14	13	15
	Texture	25	25	25	27
Microbes	Microbial community	1	1	1	1
	Microbial biomass	4	5	4	5
	Microbial activity	4	4	3	4
<b>Disturbance</b>					
Geomorphology	Permafrost	10	11	12	12
	Cryoturbation	6	8	7	8
	Fluvial activity	4	4	4	4

	Hummocks & hollows	9	10	10	10
Herbivory	Reindeer	4	5	5	5
	Muskoxen	3	3	3	3
	Small mammals	1	1	1	1
	Goose	4	4	4	4

Supplementary Material 5. Dominant species as informed by the authors. From vegetation informed only at genera-level, *Carex* (30 studies), *Sphagnum* (19 studies), *Eriophorum* (12 studies), and *Salix* (7 studies), were studied the most.

Species	Functional group	Number of studies
<i>Betula nana</i>	Deciduous shrub	29
<i>Empetrum hermaphroditum/nigrum</i>	Evergreen shrub	24
<i>Eriophorum vaginatum</i>	Graminoid	23
<i>Vaccinium uliginosum</i>	Deciduous shrub	19
<i>Vaccinium vitis-idaea</i>	Deciduous shrub	17
<i>Eriophorum angustifolium</i>	Graminoid	15
<i>Cassiope tetragona</i>	Evergreen shrub	11
<i>Rubus chamaemorus</i>	Forb	11
<i>Eriophorum scheuchzeri</i>	Graminoid	10
<i>Carex aquatilis</i>	Graminoid	9
<i>Dryas octopetala</i>	Evergreen shrub	9
<i>Carex stans</i>	Graminoid	8
<i>Rhododendron tomentosum</i>	Evergreen shrub	8
<i>Salix arctica</i>	Deciduous shrub	8
<i>Salix glauca</i>	Deciduous shrub	8
<i>Salix pulchra</i>	Deciduous shrub	8
<i>Rhododendron subarcticum</i>	Evergreen	7

	shrub	
<i>Andromeda polifolia</i>	Evergreen shrub	6
<i>Dryas integrifolia</i>	Evergreen shrub	6
<i>Dupontia psilosantha</i>	Graminoid	6
<i>Salix polaris</i>	Deciduous shrub	6

#### Supplementary Material References.

Dinerstein E, Olson D, Joshi A, et al. (2017) An Ecoregion-Based Approach to Protecting Half the Terrestrial Realm. *BioScience* 67(6): 534–545.

Fick, S.E. and R.J. Hijmans, 2017. Worldclim 2: New 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology*.

Hengl T, Mendes de Jesus J, Heuvelink GBM, et al. (2017) SoilGrids250m: Global gridded soil information based on machine learning. *PLOS ONE* 12(2): 1–40.

Hugelius, G., Bockheim, J. G., Camill, P., Elberling, B., Grosse, G., Harden, J. W., Johnson, K., Jorgenson, T., Koven, C. D., Kuhry, P., Michaelson, G., Mishra, U., Palmtag, J., Ping, C.-L., O'Donnell, J., Schirrmeister, L., Schuur, E. A. G., Sheng, Y., Smith, L. C., Strauss, J., and Yu, Z. (2013) A new data set for estimating organic carbon storage to 3 m depth in soils of the northern circumpolar permafrost region. *Earth System Science Data* 5: 393–402.

USGS (2004) Shuttle radar topography mission. Global land cover facility, University of Maryland, College Park, Maryland, February 2000.

NASA (2017). Nasa Earth Observations. Vegetation index (NDVI) 16 day – TERRA/MODIS.